




**New Jersey Department of Health and
Senior Services
Communicable Disease Service**



**Respiratory Virus Surveillance Report¹
Week ending January 7, 2012 (MMWR week 1)**

SYNOPSIS

Influenza Activity Level ²		
State Activity Week ending 1/7:		
LOW		
Current week Last year: LOW		
Regional ³ Data		
Northwest	LOW	
Northeast	LOW	
Central West	LOW	
Central East	LOW	
South	LOW	

ILI Activity ⁴				
	Percent ILI/Absenteeism			Baselines
	Current week (range by county)	Last week Current year	Current week Last year	Non-season ⁵ Season ⁶ (3 low, 3 high)
Long Term Care Facilities	0.54 (0.00-6.19)	0.41	0.71	0.64 (0.66, 0.89)
Schools (absenteeism)	5.20 (2.77-7.28)	0.00	5.68	4.32 (4.86, 5.21)
Emergency Departments	3.97 (0.00-20.40)	3.88	4.49	2.68 (3.37, 3.89)


Viral Activity ⁷			
	Current Week	Past 3 Weeks	Cumulative Total
Influenza H1N1 (2009)	0	0	0
Influenza H3N2	0	0	4
Influenza B	0	0	1
Respiratory Syncytial Virus (RSV)	264	722	2109

Report also available at:
<http://nj.gov/health/flu/fluinfo.shtml>


Virologic Surveillance⁷

NJ Public Health and Environmental Laboratory (PHEL) Influenza Positive (PCR)

Result by Week




Result by Region³




Influenza Positive (PCR) – Commercial/Acute Care Laboratories


Result by Week




Result by Region³



Influenza Rapid Antigen Result by Week





Respiratory Syncytial Virus (RSV) Results by Week





Influenza-like Illness Surveillance

Emergency Department ⁸	
Daily Visits and Admission	Visits – Tuesday Only




School Absenteeism	Long Term Care Facilities
--------------------	---------------------------



Respiratory Outbreaks in Long Term Care Facilities⁹

Cumulative outbreaks 2011-2012 season	3
No. outbreaks last 3 weeks	1
Regions with recent outbreaks	Northeast

ILINet Providers



Additional Information

A second report containing information about age specific illness, hospitalization and deaths will be produced on a monthly basis or as needed when important information needs to be disseminated.

For additional information regarding influenza surveillance please visit the following websites.

<http://nj.gov/health/flu/surveillance.shtml>

<http://www.cdc.gov/flu/>

Footnotes:

1. This report represents activity occurring in New Jersey related to influenza and RSV. In addition, reports of other circulating respiratory viruses will be included when available.
2. Activity levels for the state and region are defined in Table 1 and 2 at the end of this document.
3. The following is a breakdown of counties contained within each public health region: Northwest: Morris, Passaic, Sussex, Warren; Northeast: Bergen, Essex, Hudson; Central west: Hunterdon, Mercer, Somerset; Central East: Middlesex, Monmouth, Ocean, Union; South: Atlantic, Burlington, Camden, Cape May, Salem, Cumberland, Gloucester
4. Influenza-like illness (ILI) is defined as fever ($> 100^{\circ}\text{F}$ [37.8°C], oral or equivalent) and cough and/or sore throat (in the absence of a known cause other than influenza). For long term care facilities, fever is defined as 2° above baseline temperature.
5. Non-season baseline is calculated by taking the average of statewide percentages of ILI for a 6 year (2006, 2007, 2008, 2009, 2010, 2011) period during months when influenza is less likely to be circulating (May-August).
6. Three year seasonal averages are determined by calculating the average percent ILI/absenteeism for each influenza season (October to May). These averages are ranked and the three highest and lowest overall season averages were selected. The three highest and lowest numbers were then averaged to obtain a single high and single low value. The season which contribute to the high and low value vary by entity type and are as follows : LTCF (High: 05-06, 06-07, 07-08; Low: 08-09, 09-10, 10-11), ED (High: 05-06, 06-07, 09-10; Low: 07-08, 08-09, 10-11) and schools (High: 06-07, 07-08, 08-09 ;Low: 05-06, 09-10, 10-11). A week by week average was also calculated using the average of the seasons listed above for each entity type.
7. Viral activity: Real-time polymerase chain reaction (PCR) results are obtained from electronic laboratory transmission submitted by acute care, commercial and public health laboratories to CDRSS. Rapid influenza test data and respiratory syncytial virus data are acquired from facilities reporting rapid influenza tests via the National Respiratory and Enteric Virus Surveillance System (NREVSS) or CDRSS ILI module. Counts for cumulative totals begin with week ending October 8, 2011. Three week count data includes current week and two prior weeks. Data presented for RSV and rapid influenza testing represent information for the week prior to the current report week.
8. Daily visits and admission associated with ILI from emergency department data is collected via EpiCenter and Hippocrates. Prior to these systems, data on ILI visits were only recorded one day per week usually on Tuesday. This system is maintained as a large amount of historical data allows for better seasonal comparisons.
9. Only LTCF outbreaks reported to NJDHSS that received an outbreak number are recorded in this report.

Table 1 Influenza Activity Level – Definitions for State Activity				
<u>NJ Level</u>	<u>CSTE Level</u>	<u>Definition</u>		
		<u>ILI Activity/Outbreaks</u>		<u>Lab Activity</u>
Low	No Activity	ILI activity at or below baseline AND no detected outbreaks	AND	No lab confirmed cases
	Sporadic	Low ILI activity detected OR one lab confirmed outbreaks anywhere in the state	AND	Sporadic isolation of laboratory confirmed influenza
Moderate	Local	Increase in ILI activity OR two or more lab confirmed outbreaks in one public health region (Other regions not experiencing increased ILI activity)	AND	Recent (within 3 weeks) laboratory activity in the region with increased ILI
	Regional	Increase in ILI activity OR two or more lab confirmed outbreaks in at least 2 public health regions (Other regions not experiencing increased ILI activity)	AND	Recent (within 3 weeks) laboratory activity in the region with increased ILI
High	Widespread	Increase in ILI activity OR two or more lab confirmed outbreaks in > 2 public health regions	AND	Recent (within 3 weeks) laboratory activity in the region with increased ILI

Table 2 Influenza Activity Level – Definitions for Public Health Regions			
<u>NJ Level</u>	<u>Definition</u>		
	<u>ILI Activity/Outbreaks</u>		<u>Lab Activity</u>
Low	Low ILI activity detected OR one lab confirmed outbreaks anywhere in the region	AND	Sporadic isolation of laboratory confirmed influenza anywhere in the region
Moderate	Increased ILI activity in less than half of the counties in the region OR two lab confirmed outbreaks in the public health region	AND	Recent (within 3 weeks) laboratory activity in same counties of the region with increased ILI
High	Increased ILI activity in more than half of the counties in the region OR three or more lab confirmed outbreaks in the region	AND	Recent (within 3 weeks) laboratory activity in more than half of the counties in the region with increased ILI

Notes:

ILI activity: Systems used to detect increases in ILI activity include: ILINet (i.e., sentinel providers), school absenteeism data, ED ILI visits and admission collected via Hippocrates and EpiCenter systems, LTCF ILI data, LTCF outbreak data, and information on influenza mortality (122 city, influenza associated death report).

Lab Activity: Virologic surveillance data from PHEL will be used as the primary data source for the above levels. However, rapid influenza test data and commercial laboratory data will also be considered when determining the appropriate activity levels.



	Long Term Care			Schools			Hospital Emergency Department		
County	# Enrolled	# Reports Rec'd	% ILI	# Enrolled	# Reports Rec'd	% Absent	# Enrolled	# Reports Rec'd	% ILI
ATLANTIC	3	2	1.45	13	11	7.14	5	4	0.82
BERGEN	6	6	0.06	38	23	4.88	5	5	3.88
BURLINGTON	6	2	0.00	115	49	4.81	3	3	0.23
CAMDEN	5	0	0.00	17	0	0.00	7	1	6.90
CAPE MAY	7	4	0.58	11	9	4.81	1	0	0.00
CUMBERLAND	2	2	1.12	11	7	7.13	3	0	0.00
ESSEX	25	3	0.99	47	7	2.92	11	8	4.44
GLOUCESTER	5	3	0.34	5	5	5.12	2	1	3.03
HUDSON	16	4	0.00	86	62	7.28	6	5	5.32
HUNTERDON	4	3	0.40	8	8	2.77	1	0	0.00
MERCER	11	1	6.19	26	9	3.39	6	2	2.56
MIDDLESEX	17	11	0.25	25	13	3.41	5	5	4.83
MONMOUTH	12	2	0.00	26	11	5.09	5	4	4.22
MORRIS	4	0	0.00	5	0	0.00	4	4	3.84
OCEAN	12	2	0.00	56	22	5.20	4	1	20.40
PASSAIC	9	3	1.52	31	16	4.67	3	3	2.92
SALEM	2	1	0.00	5	2	5.89	1	1	0.00
SOMERSET	5	1	1.57	87	29	3.70	2	0	0.00
SUSSEX	5	5	0.71	23	22	4.68	2	2	0.00
UNION	3	0	0.00	195	30	4.88	5	5	1.79
WARREN	4	3	0.00	25	16	4.98	2	2	1.74
NW Region	22	11	0.92	84	54	4.74	11	11	2.88
NE Region	47	13	0.15	171	92	6.61	22	18	4.51
CW Region	20	5	2.04	121	46	3.50	9	2	2.56
CE Region	44	15	0.17	302	76	4.74	19	15	5.39
South Region	30	14	0.71	177	83	5.33	22	10	1.58
State Total	163	58	0.54	855	351	5.20	83	56	3.97

County	RSV Tests		Flu Tests		
	# Positive	Total Tests Performed	# Positive	Total Tests Performed	
ATLANTIC	11	30	0	45	
BERGEN	0	1	0	8	
BURLINGTON	17	29	0	44	
CAMDEN	16	46	0	18	
CAPE MAY	4	9	0	9	
CUMBERLAND	19	49	0	0	
ESSEX	47	111	4	74	
GLOUCESTER	11	25	0	30	
HUDSON	14	33	0	0	
HUNTERDON	12	21	0	30	
MERCER	4	13	0	5	
MIDDLESEX	57	146	0	76	
MONMOUTH	35	135	1	206	
MORRIS	4	10	0	20	
OCEAN	0	0	0	0	
PASSAIC	0	0	0	0	
SALEM	0	1	0	0	
SOMERSET	0	0	0	0	
SUSSEX	0	6	0	18	
UNION	8	33	0	5	
WARREN	5	19	0	19	
NW Region	9	35	0	57	
NE Region	61	145	4	82	
CW Region	16	34	0	35	
CE Region	100	314	1	287	
South Region	78	189	0	146	
State Total	264	717	5	607	