


Cardiac Surgery

in New Jersey, 2013



Health Care Quality Assessment
Office of Population Health

November, 2016



Chris Christie
Governor
Kim Guadagno
Lt. Governor



Cathleen D. Bennett
Commissioner



Message from the Commissioner

I am pleased to present the 17th Cardiac Surgery Report in New Jersey, the state's consumer report on coronary artery bypass graft (CABG) surgery. This report summarizes mortality, length of stay and infections among patients who underwent bypass surgery in New Jersey hospitals in 2013.

New Jersey's cardiac bypass surgery operative mortality rate has declined by 63.1 percent between 1994 and 2013. This is a tribute to the continued commitment of our state's cardiac surgery centers and surgeons to making cardiac bypass surgery safer.

The Department of Health continues to work closely with the Cardiovascular Health Advisory Panel (CHAP) to bring consumers and providers the best possible data on cardiac bypass surgery to assess outcomes. In addition to CABG surgery mortality associated with hospitals, this report also includes the number of cardiac surgeries physicians performed, their surgery outcomes and summary of statewide cardiac bypass surgery-related infections.

I would like to thank the CHAP members for their important efforts in supporting quality improvement in cardiac services in New Jersey.

In facing cardiac bypass surgery, patients and their families have questions and concerns. We hope this report answers some of those questions and helps patients discuss concerns and treatment options with their physicians.



Cathleen D. Bennett
Commissioner

Table of Contents

Executive Summary iii

Introduction 1

How to Use this Report 1

Cardiovascular Health Advisory Panel 2

Heart Disease and Cardiac Surgery in New Jersey 2

Treatment Options 2

Definition of Operative Mortality 2

Performance Data 3

Risk-adjusted Mortality 3

Performance Reports Lead to Improvement 3

Hospitals 4

Surgeons 4

Volume Affects Quality 4

Bypass Surgery Volume at New Jersey Hospitals in 2013 4

Hospital Risk-Adjusted Mortality 4

Statewide Bypass Surgery Related Infections 8

Length of Stay by Hospital 9

Individual Surgeon Performance 10

Statewide Trends in Risk-Adjusted Bypass Surgery Mortality Rates:

Pooled Estimates 18

Figure 1: Number of Isolated Coronary Artery Bypass Graft Surgeries vs. Other Cardiac Surgeries (2013) 6

Figure 2: Risk-Adjusted Operative Mortality Rate by Hospital (2013) 7

Figure 3: Risk-Adjusted Operative Mortality and Length of Stay by Hospital, 2013. 9

Figure 4: Surgeon Risk-Adjusted Operative Mortality Rate (2012-2013) 11

Figure 5: Trends in Statewide Bypass Surgery Mortality Rates 18

Table 1: Statewide In-hospital Infection Rate and Operative Mortality Rate by Infection Type, 2013. 8

Table 2: Risk-Adjusted Mortality Rate and Post-Surgery Length of Stay by Surgeon (2012-2013) 15

Appendix A: Questions and Answers 19

Appendix B: New Jersey’s Cardiovascular Health Advisory Panel Members and DOH Cardiac Surgery Report Team 20

Appendix C: Statewide Observed In-hospital and Operative Mortality Rates 21

Appendix D: Summary of Methods Used in this Report 22

References 31

Executive Summary

This report is for patients and families considering coronary artery bypass graft (CABG) surgery. It summarizes the results of a study of CABG surgery in New Jersey and answers many of the questions you may have about this common procedure.

An important goal of this report is to give you, the patient, and your family information that will help you have more informed discussions with your physicians. Since every patient has different health concerns and risks, we encourage you to discuss the information in this report with your physicians, who can answer your questions and concerns.

Another important goal of this report is to give New Jersey hospitals and surgeons meaningful data they can use in assessing quality of care related to CABG surgery. There is strong evidence, from the handful of states with similar studies, that this kind of information prompts hospitals to examine their process of care in order to improve the overall quality of CABG surgery, prevent infections, and ultimately save lives.

For this study, the Department of Health (Department) collected data on 8,251 patients undergoing open heart surgery at 18 hospitals in 2013. Of these patients, 3,881 had CABG surgery with no other major surgery during the same admission, i.e. isolated CABG surgery (or simply referred to as bypass surgery in this report).

This study was a collaborative effort with a select committee of experts known as the Cardiovascular Health Advisory Panel (CHAP), which includes physicians who specialize in cardiac surgery, cardiologists and other health care professionals.

How to Use This Report

Hospitals and doctors are not the same in their specialties and expertise. Some are better equipped than others to handle patients with different health conditions. These differences will influence the quality of care you receive and the outcomes of your bypass surgery.

Many consumers want a doctor's recommendation on hospitals and surgeons. Frequently, people collect as much information as possible to make informed decisions. This report will provide some of that information.

However, this report is not intended to be used alone. It is designed to provide important information to help you make informed decisions. There are many factors to consider in determining the best hospital for you. Among these are your own personal health risks as well as the experience certain hospitals have treating patients with those risk factors. Before you make your decisions, you should discuss this report with the physician, usually a cardiologist, who refers you for cardiac surgery. The cardiologist's knowledge and expertise will be a valuable guide in making your decision.

Key Findings

The Department analyzed the bypass surgery data using a statistical method to assess hospital and surgeon performance. Before analyzing the data, the Department performed extensive error checks on the entire open heart surgery data, sampled medical records from each hospital for independent medical audit and consulted with the clinical panel of the CHAP. The statistical analysis took into account the patient's health status before surgery as well as demographic factors. This process is commonly known as "risk-adjustment" and allows for fair comparisons among hospitals and surgeons treating diverse patient populations.

Some key findings of the 2013 data analysis are as follows:

Statewide Summary

- 47.0 percent (3,881) of the 8,251 total open heart surgeries performed in New Jersey in 2013 were bypass surgeries.
- Of the 3,881 bypass surgery patients, 61 died while in the hospital or within 30 days after surgery. The statewide observed operative mortality rate for bypass surgery patients in 2013 was 1.57 percent.

- When comparing 2012 and 2013 on a risk-adjusted basis, the mortality rate decreased 16.6 percent.
- A review of the 20 years of pooled data suggests that the risk-adjusted bypass mortality rate in New Jersey declined 63.1 percent between 1994 (4.37%) and 2013 (1.61%).

Mortality Rate by Hospital and by Surgeon

- Despite the variations in bypass mortality rates among hospitals and surgeons, the quality of care delivered by most hospitals and surgeons were similar to the statewide performance.
- In 2013, one hospital, St. Francis Medical Center, had a statistically significantly higher risk-adjusted mortality rate than the statewide rate.
- No hospital had statistically significantly lower risk-adjusted mortality rate than the statewide rate. Although their rates were not statistically significantly different from the statewide rate, it is nevertheless notable that Englewood Hospital and University Hospital had no bypass surgery deaths in 2013.
- In the period 2012-2013, one surgeon had a statistically significantly higher risk-adjusted mortality rate than the statewide rate.
- No surgeon had a statistically significantly lower risk-adjusted mortality rate than the statewide rate in the period 2012-2013.
- Although their rates were not statistically significantly different from the statewide rate, it is nevertheless notable that a few surgeons, including some who performed less than 100 bypass surgeries, had no bypass surgery death during the two-year period. Among surgeons who performed 100 or more bypass surgeries during the period 2012-2013, Dr. James Klein from Englewood Hospital and Medical Center had no bypass surgery death.

Pre-surgery Patient Risk Factors

- Key factors that are associated with a patient's chance of surviving the operation include*:
 - ❖ patient's age;
 - ❖ whether the patient had various preoperative risk factors, such as certain types of cerebrovascular disease, immunosuppressive therapy or peripheral vascular disease;
 - ❖ whether the patient had preoperative cardiac status such as cardiogenic shock, congestive heart failure, or symptoms of severe cardiac insufficiency.

Post-surgery Length of Stay

- The average post-surgery length of hospital stay for a typical bypass surgery patient in 2013 was 7.16 days, which was shorter than that of 7.42 days in 2012.
- The risk-adjusted length of stay by hospital ranged from 4.56 days at St. Joseph's Regional Medical Center to 7.60 days at Hackensack University Medical Center in 2013.
- There were also differences in length of stay by surgeon. Risk-adjusted average length of stay by individual eligible surgeon in the period 2012-2013 ranged from 4.47 days to 8.14 days.

Post-surgery Infections

- In 2013, 5.28 percent of patients had some type of infection, including pneumonia, following bypass surgery. The overall infection rate decreased by 21.8 percent from 6.75 percent in 2012 to 5.28 percent in 2013 (not risk-adjusted).
- As expected, bypass surgery patients who developed infections after surgery had a much higher mortality rate (9.27 percent vs. 1.14 percent) and a longer hospital stay compared to those who had no infections (17.79 days vs. 6.57 days).

* *More information on risk factors and methods used in this report are presented in Appendix D.*

Introduction

This report is for patients and families of patients facing the possibility of coronary artery bypass graft (CABG) surgery. It provides mortality rates for the 18 hospitals that performed cardiac surgery in 2012 and 2013 and the physicians performing this procedure in 2012 and 2013. As part of the Department's continued effort to provide information to consumers, this report includes information on hospital length of stay and infections following CABG surgeries. The report provides risk-adjusted length of hospital stay after CABG surgery by hospital and by eligible surgeon (i.e., surgeon who performed at least 100 isolated CABG operations in one hospital in the years 2012 and 2013 combined). The rates of infections are reported for the state as a whole.

An important goal of the report is to give you, the patient, and your family information that will help you have more informed discussions with your physician. Since every patient has different health concerns and risks, we encourage you to discuss the information in this report with your physician, who can best answer your questions and concerns.

Another important goal of this analysis is to give hospitals data they can use in assessing quality of care related to CABG surgery. There is strong evidence, from other states with similar reports, that this information encourages hospitals to examine their processes of care and make changes that can improve quality of care, prevent infections, and ultimately save lives.

For this report, the Department of Health collected data on 3,881 patients who had CABG surgery with no other major surgery during the same admission (simply referred to as isolated CABG surgery or bypass surgery in this report) in 2013. These are the most recent years for which death certificate data used to calculate mortality up to 30 days after discharge are

available. The data have been "risk-adjusted," which means that they were adjusted to take into account the patient's health conditions before surgery. The risk-adjustment process allows for fair comparisons among hospitals and surgeons treating diverse patient populations.

New Jersey's mortality rate for bypass surgery has shown marked decline since public reporting began with 1994 data. The risk-adjusted mortality rate has decreased 63.1 percent from 4.37 percent to 1.61 percent between 1994 and 2013, which is statistically significant. A difference is called "**statistically significant**" when it is too large to be due to chance or random variation.

The observed mortality rate in 2013 was 1.57 percent, which was lower than the 2012 mortality rate of 2.01 percent. The risk-adjusted mortality rate decreased 16.6 percent between 2012 and 2013, which is not statistically significant (Appendix D).

How to Use this Report

Hospitals and doctors are not the same in their specialties and expertise. Some are better equipped than others to handle patients with different health conditions. These differences will influence the quality of care you receive and the outcomes of your bypass surgery.

Many consumers want a doctor's recommendation on hospitals and surgeons. Frequently, people collect as much information as possible to make informed decisions. This report will provide some of that information. However, this report is not intended to be used alone. Volume, mortality rate and length of stay in this report are just some of the important factors to consider in deciding where to have cardiac surgery. There are many factors to consider in determining the best hospital for you. Among these are your own personal health risks as well as the experience certain hospitals have treating patients with those risk factors. Before you make your decisions, you should discuss this report with your physician, usually

a cardiologist, who refers you for cardiac surgery. You and your physician together can make the best choice after full consideration of your medical needs.

Cardiovascular Health Advisory Panel

A Cardiovascular Health Advisory Panel (CHAP) was established by the Commissioner of Health by Executive Order (No. 187 (2001) and amended by Executive Order 207) to provide the Commissioner with expert advice on sound cardiovascular health policy. CHAP provides advice on cardiovascular health promotion, disease prevention, standards of care, emerging technologies and their applications to cardiac services in the State, and review of the State's cardiac data for quality assessment, performance evaluation and research. CHAP's membership includes surgeons, cardiologists, nurses and professional associations and consumer representatives (See Appendix B).

Heart Disease and Cardiac Surgery in New Jersey

Heart diseases continue to be the leading causes of death of Americans with 611,105 deaths in 2013. Almost every 30 seconds, someone in the United States will suffer a heart attack, and about once every minute, someone will die from one. In New Jersey, heart diseases are the leading causes of death, accounting for 18,460 deaths in 2013. The age-standardized death rate in 2013 was 170.1 per 100,000, which was slightly higher than the national age-standardized rate of 169.8 per 100,000. (http://www.cdc.gov/nchs/data/nvsr/nvsr64/nvsr64_02.pdf, page 87, table 19).

The most common form of heart disease is coronary artery disease. Coronary artery disease occurs when the coronary arteries, which carry blood to the heart muscle, become clogged or partially blocked by fatty deposits on the artery walls. This can lead to chest pain, or angina, which is a warning sign for a heart attack. A heart attack occurs when a coronary artery is totally blocked.

Treatment Options

Treatment for coronary artery disease will vary for different patients. The choice of treatment depends on the nature and severity of the disease and other factors unique to each patient.

For some patients, lifestyle changes such as quitting smoking, eating a low-fat diet, and getting more exercise may be enough. Some patients require special medications. Others may need medical procedures such as percutaneous coronary intervention (PCI, commonly known as angioplasty) or CABG surgery. Angioplasty reduces obstructions of fatty deposits in coronary arteries and has become an increasingly common treatment method. CABG surgery uses an artery or vein taken from another part of the body to divert blood around the clogged part of a patient's artery or arteries.

This report is about coronary artery bypass graft (CABG or bypass) surgery outcomes. It describes the performance records of 18 hospitals in New Jersey that offered this type of surgery in 2012 and 2013, as well as the surgeons who performed this operation at least 100 times between January 2012 and December 2013 in a hospital.

Definition of Operative Mortality

Beginning with the 2000 report¹, the Department, after consulting with the CHAP, included in its definition of "operative mortality" deaths up to 30 days post-surgery or deaths occurring during the hospital stay in which the surgery was performed, no matter how many days after the procedure. Deaths occurring within 30 days after surgery, but post-discharge, have been identified by matching patient records in the Department's Open Heart Surgery database against the State's official death records.

¹ Prior to 2000, the Department defined patient death for this report as in-hospital death before discharge from the hospital after bypass surgery. As a result, patients who died after being discharged home or to post-acute care facilities were not counted for purposes of calculating bypass surgery mortality rates. This caused concerns about "gaming" of outcomes through discharge practices.

Further, in an attempt to continuously improve the quality of data used in assessing bypass surgery mortality, the Department, in consultation with CHAP, reviewed the way operative procedures are coded for the purpose of the cardiac surgery report in New Jersey. The Department issued an operative procedure coding guide to be followed by all hospitals starting with 2005 data. This guideline was designed to avoid differential reporting of operative procedures by hospitals.

Applying the revised definitions of mortality, the Department also recalculated the statewide bypass surgery mortality rates for the prior years, in order to analyze the trend over time. Trend in operative mortality rate estimates from 1994 to 2013 are presented in Figure 5. Appendix D, Table D3 also presents the statewide operative mortality rate estimates for the period 1994-2013.

Performance Data

In an isolated CABG (bypass) surgery, no other major heart procedure is performed at the same time. In 2013, the number of people who died during the hospitalization in which the operation was performed, or after discharge but within 30 days of the surgery, was 61. This represents 1.57 percent of the 3,881 who had bypass surgery in 2013. This rate is referred to as statewide operative mortality rate. This statewide operative mortality rate (1.57 percent) is used as the yard stick in evaluating hospital performance.

Risk-Adjusted Mortality

In evaluating the performance of hospitals and individual surgeons, it would be unfair to make comparisons only on the basis of how many patients died. The mortality risk for patients undergoing bypass surgery varies significantly with how healthy patients are prior to surgery. For instance, an 85-year-old who had a certain type of cerebrovascular disease and was in cardiogenic shock at the time of surgery would be at higher risk during this surgery than a 50-year-old who had no history of chronic disease.

In order to produce fair comparisons, the Department applied a method that estimates **risk-adjusted mortality rates**. Each hospital was required to submit data which contain a risk profile for each patient undergoing bypass surgery. The risk-adjusted mortality rate assigns “extra credit” to hospitals and surgeons with sicker patient populations, in order not to disadvantage them in the performance comparisons.

Key factors that are associated with a patient’s chance of surviving the bypass operation include:

- patient’s age;
- whether the patient had various preoperative risk factors, such as certain types of cerebrovascular disease, immunosuppressive therapy or peripheral vascular disease;
- whether the patient had preoperative cardiac status such as cardiogenic shock, congestive heart failure, or symptoms of severe cardiac insufficiency.

Weights derived from the statistical model were assigned for each key risk factor and **risk-adjusted mortality rates** were calculated for each hospital as fair basis for comparison (see Appendix D for more details).

Performance Reports Lead to Improvement

This performance report is for use not only by you and your doctors, but also by hospitals to improve the quality of their care and their patients’ outcomes. On a risk-adjusted basis, the New Jersey statewide risk-adjusted mortality rate for bypass surgery declined 63.1 percent from 4.37 percent in 1994 to 1.61 percent in 2013 (see Appendix D, Table D3). Evidence both from New Jersey and other states that have published similar performance reports (i.e. California, Massachusetts, New York and Pennsylvania) suggests that these reports contribute to the decline in mortality rates and improve the overall quality of bypass surgery.

Hospitals

This report provides risk-adjusted mortality rates for each of the 18 hospitals in New Jersey that were licensed to perform coronary artery bypass graft surgery in 2013. You will see that there are substantial variations among the 18 cardiac surgery hospitals. Through statistical analysis, the Department is able to determine in which cases the variations reflect real differences in performance after accounting for levels of risk among patients.

Nevertheless, these data should not be used as the sole factor in making choices about hospitals, but should be part of the discussion between you and your doctor.

Surgeons

A risk-adjusted mortality rate was also calculated for each of the 30 surgeons who performed at least 100 bypass operations in one hospital in the years 2012 and 2013 combined. Even though two years of data were combined, several surgeons still fell short of the 100 cases the Department considers the minimum needed to calculate reliable risk-adjusted mortality rates. The Department recognizes that the volumes of some surgeons may be low because they had left those facilities during the year. Statistics for these low-volume surgeons are grouped under the hospital where the operations took place, in a category called **“All Others.”** These surgeons are listed by name but with no risk-adjusted mortality rates, since their small numbers do not permit an accurate indication of their performance (Table 2). This report shows the total number of open heart and bypass surgeries these low volume surgeons performed, as well as their number of bypass surgery operative deaths.

Volume Affects Quality

Many studies nationally and in other states have shown that, in general, hospitals and surgeons that perform bypass surgery more

frequently have lower patient mortality rates. New Jersey’s data also confirm this general trend. However, there are exceptions, and a number of hospitals with low volumes have results that are in line with the statewide rate.

Bypass Surgery Volume at New Jersey Hospitals in 2013

Bypass surgery is the most common type of cardiac surgery accounting for 47.0 percent in 2013. Figure 1 shows the number of bypass operations performed in 2013 in each of the 18 hospitals. You can see that some hospitals do more of these procedures than others, with bypass volumes ranging from a low of 29 at University Hospital to a high of 643 at Morristown Medical Center. Bypass surgery volume has been declining in New Jersey starting in 2000 while angioplasty has stabilized at a higher level. Between 2000 and 2013, the number of bypass surgeries in New Jersey has declined by 52.8 percent.

Hospital Risk-Adjusted Mortality

Figure 2 shows the risk-adjusted mortality rate for each New Jersey hospital performing bypass surgery in 2013². The risk-adjusted mortality rate takes into account the patient’s risk factors before surgery as well as the actual mortality rate after the surgery, in order to make a fair assessment of hospital performance.

In trying to determine hospital or surgeon performance, it is important to account for the fact that some differences occur simply due to chance or random variation. Statistical tests are performed on the risk-adjusted bypass mortality estimates so that we can be as certain as possible that the differences are due to actual variations in performance. A difference is called **“statistically significant”** when it is too large to attribute to chance or random variation.

Each hospital's and each surgeon's mortality rate reflects three components: *the quality of*

² These data may not reflect current performance of a specific hospital, which may have revamped its program since then.

their care, the patient's risk factors that affect mortality, and an element of random variation. Readers of this report should be interested only in the first component, the quality of care delivered by hospitals and surgeons. We use a nationally-accepted risk-adjustment method to control for the second component, risk factors of bypass surgery patients seen by hospitals and surgeons. Because the third component, *random variation*, cannot be observed to be controlled for in the statistical model, we estimate how much higher or lower the risk-adjusted mortality rate could have been given the impact of random variation, using a confidence interval given at the 95% level.

In Figure 2, the dark line in the middle of each hospital's bar represents its estimated risk-adjusted mortality rate. When estimating rates using data, however, we cannot be sure if this number is the actual rate for the facility and not due to chance. We can only be relatively sure that the true rate falls somewhere within the bar. In analyzing data, we use what is called a "95 percent confidence interval," and the bar represents the lower and upper limits of this confidence interval. We are 95 percent confident that the hospital's actual risk-adjusted mortality rate falls within the range shown by the bar. Another way of saying it is that the bar represents the statistical margin of error for the estimation of that rate.


The vertical line on Figure 2 represents New Jersey's statewide bypass surgery operative mortality rate per 100 cases for 2013, i.e. 1.57. Each hospital's performance is displayed graphically in relation to this statewide rate.

Figure 2 indicates 17 hospitals have bars that cross the statewide mortality rate line (1.57 percent). That means that their risk-adjusted mortality rates were not statistically different from the statewide rate. St. Francis Medical Center has its bar completely to the right of the statewide rate indicating that this hospital had a statistically higher risk-adjusted mortality rate than the statewide rate.

When using this report, it is important to remember that the charts are designed to show whether a hospital's or surgeon's risk-adjusted mortality rate is significantly above or below the statewide rate, or whether a rate is statistically the same as the statewide rate. Thus, it is more important to view the bars in relation to the statewide mortality rate line than it is to examine the individual calculated rates on the bars. The chart should not be used to make hospital-to-hospital or surgeon-to-surgeon comparisons, only to compare hospitals and surgeons to the statewide rate.


In examining the charts, you will see that some bars are shorter than others. The bar is shorter for hospitals or surgeons performing more surgeries, and longer for those with lower volumes. This reflects the fact that larger numbers -- in this case, more surgeries -- increase the precision of a statistic.

Figure 1
Number of Isolated Coronary Bypass Graft Surgeries vs. Other Open Heart Surgeries, 2013



Source: New Jersey Department of Health

Figure 2
Risk-Adjusted Operative Mortality Rate* by Hospital (2013)



SOURCE: New Jersey Department of Health

* = Operative Mortality includes: (1) all deaths occurring during the hospitalization in which the operation was performed, even after 30 days; and (2) those deaths occurring after discharge from the hospital, but within 30 days of the procedures.

(+) = Risk-adjusted mortality rate significantly higher than the New Jersey statewide mortality rate based on 95 percent confidence interval.

Statewide Bypass Surgery Related Infections

The Department has included information on bypass surgery in-hospital infections as an additional tool to monitor hospital performance. The statewide infection rates are provided as one more factor to be considered by policy makers and others involved in quality of care monitoring.

Infections reported in the Open Heart Surgery database included sternal-deep infection (involving muscle, bone and/or mediastinum requiring operative intervention), thoracotomy, leg infections, septicemia (presence of bacteria in the blood stream) and urinary tract infections (UTI). The table also includes post-operative pneumonia. For comparison purposes, statewide infection rates, the corresponding mortality rates and the average length of stay are presented in Table 1 to provide perspective to the statewide rates.

Table 1 shows that, statewide, 5.28 percent of patients who underwent bypass surgery had some type of infection (including pneumonia). 2.65 percent of

bypass patients had pneumonia, 2.04 percent of patients had UTI and 0.49 percent had septicemia.

Observed bypass surgery mortality for those who had infections (9.27%) was more than eight times as high as those who did not (1.14%). In addition, patients who developed post-surgery infections stayed in the hospital more than two and half times as long (17.79 days) as those who had no infection (6.57 days).

Septicemia had the highest mortality rate of 26.32 percent among all infections reported, followed by pneumonia (14.56%), and UTI (6.33%).

Statewide, overall infection rate after bypass surgery decreased by 27.8 percent from 6.75 percent in 2012 to 5.28 percent in 2013 (not risk-adjusted). The decline in infection rate occurred to most infections reported, from leg infection (0.51% to 0.33%) to septicemia (0.91% to 0.49%), UTI (2.81% to 2.04%) and pneumonia (3.48% to 2.65%). Infections from sternal-deep (0.35% to 0.41%) and thoracotomy (0.00% to 0.03%) increased slightly.

Table 1
Statewide In-hospital Infection Rate and Operative Mortality Rate by Infection Type, 2013

	Number of Cases	Infection Rate (%)	Operative Mortality*		Average Length of Stay (in Days)
			Number	Rate (%) (Observed)	
Cases with Infections	205	5.28	19	9.27	17.79
Sternal-Deep	16	0.41	0	0.00	18.56
Thoracotomy	1	0.03	0	0.00	34.00
Leg	13	0.33	0	0.00	23.62
Septicemia	19	0.49	5	26.32	32.47
UTI	79	2.04	5	6.33	14.96
Pneumonia	103	2.65	15	14.56	18.20
Cases without Infections	3,676		42	1.14	6.57
Total CABG cases	3,881		61	1.57	7.16

SOURCE: New Jersey Department of Health

* Operative Mortality includes:

- (1) all deaths occurring during the hospitalization in which the operation was performed, even after 30 days; and
- (2) those deaths occurring after discharge from the hospital, but within 30 days of the procedures.

Length of Stay by Hospital

The Department has included information on post-surgery length of stay as an additional tool to monitor hospital and surgeon performance on bypass surgery. The statewide post-surgery length of stay is 7.16 days.

The risk-adjustment length of stay model excludes in-hospital deaths, very low lengths of stay (low outliers) and very long lengths of stay (high outliers) while fitting the regression model to reduce outlier effects on the model.


The risk-adjusted length of stay by hospital are displayed in Figure 3 and compared against their respective risk-adjusted mortality rates.

Figure 3 shows that there is a marked variation in risk-adjusted length of stay by hospital. The risk-adjusted length of stay by hospital ranges from a low of 4.56 days at St. Joseph’s Regional Medical Center to a high of 7.60 days at Hackensack University Medical Center. The correlation between hospital mortality rate and length of stay is not statistically significant.

Length of stay data for individual surgeons are presented later in this report.

Figure 3
Risk-Adjusted Operative Mortality and Length of Stay by Hospital, 2013

Risk-Adjusted Operative Mortality and Length of Stay by Hospital, 2013



Individual Surgeon Performance

Figure 4 and Table 2 show the risk-adjusted mortality rate for each of the 30 surgeons who performed at least 100 bypass surgery operations in one hospital in New Jersey in the years 2012 and 2013 combined³. In addition, Table 2 shows the risk-adjusted length of stay for each surgeon.

Table 2 lists surgeons by name under the hospital in which they practiced. At the end of each list of named surgeons, some hospitals have an “All Others” category. “All Others” includes all surgeons who performed too few procedures in that hospital for an individual risk-adjusted mortality rate to be calculated. Mortality rate for the “All Others” category is displayed in Table 2 only when it includes at least two or more surgeons and 25 or more bypass patients. Similarly, Figure 4 displays a bar for a surgeon only if 100 or more bypass surgeries were performed by the surgeon in one hospital in the years 2012 and 2013 combined. For a group of surgeons (i.e. All Others) a bar is shown when the group includes at least two or more surgeons and 25 or more total patients. It is important to note that some surgeons may no longer be practicing cardiac surgery in the facilities where they are listed.

Once again, the vertical line in Figure 4 represents the statewide operative mortality rate for 2012 and 2013 combined. Note that, when two years’ data are combined, the statewide operative mortality rate was 1.79 percent. If a surgeon has a bar completely to the left of the statewide line, i.e. 1.79, it means that the surgeon’s mortality rate was statistically significantly lower than the statewide rate. No surgeon had a statistically significantly lower risk-adjusted mortality rate than the statewide rate.

As is the case for some in this report, it is possible for a surgeon to have no patient deaths


and still have his/her bar cross the statewide line. Though not intuitive, this happens because the bar is the result of an upper and lower bound which includes standard errors of the estimated mortality rate. Although their rates were not statistically significantly different from the statewide rate, it is nevertheless notable that a few surgeons, including some who performed less than 100 bypass surgeries, had no bypass surgery death during the two-year period. Among surgeons who performed 100 or more bypass surgeries during the period 2012-2013, Dr. James Klein from Englewood Hospital and Medical Center had no bypass surgery death.

If a surgeon has a bar completely to the right of the statewide mortality rate line, it means that the surgeon's mortality rate was statistically significantly higher than the statewide rate for this two-year period. In 2012-2013, one surgeon, Dr. George Batsides from Robert Wood Johnson University Hospital, had a statistically significantly higher risk-adjusted mortality rate than the statewide rate.

In addition to risk-adjusted mortality for surgeons, Table 2 also shows risk-adjusted patient length of stay for each surgeon who performed at least 100 bypass surgeries in the 2012-2013 reporting period. There is marked variation in length of stay among eligible surgeons where the shortest length of stay was 4.47 days and the longest was 8.14 days. The reasons behind the wide variation in lengths of stay are not clear and need further study.

³ These data may not reflect the current performance of a specific surgeon, who may have improved his/her performance since then. Also, some surgeons listed in the cardiac surgery centers may have already left the facility since the data were reported.

Figure 4
Surgeon Risk-Adjusted Operative Mortality* Rate (2012 - 2013)




SOURCE: New Jersey Department of Health

* = Operative Mortality includes: (1) all deaths occurring during the hospitalization in which the operation was performed, even after 30 days; and (2) those deaths occurring after discharge from the hospital, but within 30 days of the procedures.

(+) = Risk-adjusted mortality rate significantly higher than the New Jersey statewide mortality rate based on 95 percent confidence interval.

** = Surgeon not currently performing CABG surgery in this hospital.

Figure 4 (continued)
Surgeon Risk-Adjusted Operative Mortality* Rate (2012 - 2013)




SOURCE: New Jersey Department of Health

* = Operative Mortality includes: (1) all deaths occurring during the hospitalization in which the operation was performed, even after 30 days; and (2) those deaths occurring after discharge from the hospital, but within 30 days of the procedures.

(+) = Risk-adjusted mortality rate significantly higher than the New Jersey statewide mortality rate based on 95 percent confidence interval.

** = Surgeon not currently performing CABG surgery in this hospital.

Figure 4 (continued)
Surgeon Risk-Adjusted Operative Mortality* Rate (2012 - 2013)




SOURCE: New Jersey Department of Health

* = Operative Mortality includes: (1) all deaths occurring during the hospitalization in which the operation was performed, even after 30 days; and (2) those deaths occurring after discharge from the hospital, but within 30 days of the procedures.

(+) = Risk-adjusted mortality rate significantly higher than the New Jersey statewide mortality rate based on 95 percent confidence interval.

** = Surgeon not currently performing CABG surgery in this hospital.

Figure 4 (continued)
Surgeon Risk-Adjusted Operative Mortality* Rate (2012 - 2013)



SOURCE: New Jersey Department of Health

* = Operative Mortality includes: (1) all deaths occurring during the hospitalization in which the operation was performed, even after 30 days; and (2) those deaths occurring after discharge from the hospital, but within 30 days of the procedures.

(+) = Risk-adjusted mortality rate significantly higher than the New Jersey statewide mortality rate based on 95 percent confidence interval.

** = Surgeon not currently performing CABG surgery in this hospital.

Table 2

Risk-Adjusted Operative Mortality* Rate and Post-Surgery Length of Stay by Surgeon (2012-2013)

Hospital and Surgeon	Total Open Heart Procedures	Number of Isolated CABG Operations	Patient Operative Deaths*	Observed Patient Mortality(%)	Expected Patient Mortality(%)	Risk-Adjusted Patient Mortality (%)	95% Confidence Interval	Risk-Adjusted Post-Surgery Length of Stay
AtlantiCare Regional Medical Center								
Axelrod, Howard	251	146	2	1.37	1.72	1.43	(0.16, 5.15)	6.08
Dralle, James, G.	240	131	1	0.76	1.68	0.81	(0.01, 4.51)	6.51
Cooper Hospital University Medical Center								
Highbloom, Richard	337	273	3	1.10	2.03	0.97	(0.19, 2.83)	5.02
All Others (13)	750	79	5	6.33	1.75	6.47	HI (2.09, 15.10)	5.95
Bowen, Frank	373	40	4					
Caputo, Francis	4	0	0					
Carpenter, Jeffrey	1	0	0					
Chovanes, John	4	0	0					
Fox, Nicole	2	0	0					
Green, Raymond	4	0	0					
Lombardi, Joseph	3	0	0					
Manis, George**	1	0	0					
Rosenbloom, Michael	344	39	1					
Seamon, Mark**	3	0	0					
Sixta, Sherry**	6	0	0					
Tsiotsias, George**	1	0	0					
Warta, Melissa**	4	0	0					
Deborah Heart and Lung Center								
McGrath, Lynn, B.	354	137	2	1.46	2.06	1.27	(0.14, 4.57)	6.26
All Others (1)	204	87	2					
Ng, Arthur, F.**	204	87	2					
Englewood Hospital & Medical Center								
Klein, James, J.	331	130	0	0.00	2.00	0.00	(0.00, 2.52)	6.35
All Others (4)	285	159	0	0.00	1.87	0.00	(0.00, 2.20)	6.32
Arnofsky, Adam	181	84	0					
Elmann, Elie, M.	27	20	0					
Praeger, Peter, I.**	50	36	0					
Somberg, Eric, D.**	27	19	0					
Hackensack University Medical Center								
Elmann, Elie, M.	441	265	2	0.75	2.20	0.61	(0.07, 2.21)	7.88
Somberg, Eric, D.	190	144	1	0.69	1.65	0.75	(0.01, 4.18)	8.14
All Others (4)	590	148	1	0.68	1.97	0.61	(0.01, 3.41)	6.55
Diluozzo, Gabriele	67	25	1					
Lee, Leonard**	61	13	0					
McCullough, Jock, N.**	435	93	0					
Praeger, Peter, I.**	27	17	0					

* = Operative Mortality includes: (1) all deaths occurring during the hospitalization in which the operation was performed, even after 30 days; and (2) those deaths occurring after discharge from the hospital, but within 30 days of the procedures.

** = Surgeon not currently performing CABG surgery in this hospital.

HI = Risk-adjusted mortality rate significantly higher than the New Jersey statewide mortality rate based on 95 percent confidence interval.

Table 2 (continued)

Risk-Adjusted Operative Mortality* Rate and Post-Surgery Length of Stay by Surgeon (2012-2013)

Hospital and Surgeon	Total Open Heart Procedures	Number of Isolated CABG Operations	Patient Operative Deaths*	Observed Patient Mortality(%)	Expected Patient Mortality(%)	Risk-Adjusted Patient Mortality (%)	95% Confidence Interval	Risk-Adjusted Post-Surgery Length of Stay
Jersey City Medical Center								
Krause, Tyrone	352	214	7	3.27	1.85	3.17	(1.27, 6.52)	5.27
All Others (2)**	103	66	0	0.00	1.17	0.00	(0.00, 8.48)	6.35
Hanhan, Ziad, G.**	54	35	0					
Predergrast, Thomas**	49	31	0					
Jersey Shore University Medical Center								
Dejene, Brook, A.	408	190	1	0.53	1.97	0.48	(0.01, 2.65)	6.34
Greeley, Drew, P.**	399	232	3	1.29	1.58	1.46	(0.29, 4.27)	6.41
Johnson, David, L.	394	194	6	3.09	1.61	3.44	(1.26, 7.49)	6.58
All Others (2)	384	172	5	2.91	1.65	3.14	(1.01, 7.33)	7.37
Neibart, Richard, M.	203	83	4					
Youdelman, Benjamin	181	89	1					
Morristown Medical Center								
Brown III, John, M.	748	185	3	1.62	1.75	1.66	(0.33, 4.84)	5.94
Magovern, Christopher, J.	648	364	3	0.82	1.86	0.79	(0.16, 2.31)	6.58
Slater, James, P.	607	365	4	1.10	1.98	0.99	(0.27, 2.53)	6.35
Xydas, Steve	620	306	6	1.96	1.76	1.99	(0.73, 4.34)	6.64
Newark Beth Israel Medical Center								
Karanam, Ravindra, N.	279	126	6	4.76	2.15	3.96	(1.45, 8.62)	7.33
Saunders, Craig, R.	341	114	1	0.88	1.26	1.24	(0.02, 6.92)	7.50
All Others (6)	548	116	2	1.72	1.59	1.94	(0.22, 7.00)	7.09
Burns, Paul, G.**	110	12	0					
Camacho, Margarita	195	17	1					
Malhotra, Sunil**	1	0	0					
Russo, Mark	150	42	0					
Sardari, Frederic, F.	36	17	1					
Simsir, Sinan**	56	28	0					
Our Lady of Lourdes Medical Center								
Kuchler, Joseph, A.	309	212	4	1.89	2.45	1.37	(0.37, 3.51)	6.05
Luciano, Pasquale	287	247	2	0.81	1.91	0.76	(0.08, 2.73)	5.72
Martella, Arthur	471	251	7	2.79	1.89	2.64	(1.06, 5.44)	4.80
All Others (2)	61	29	1	3.45	2.63	2.34	(0.03, 13.01)	5.54
Curiale, Steven**	25	21	1					
Davis, Paul**	36	8	0					
Robert Wood Johnson University Hospital								
Batsides, George, P.	555	293	13	4.44	2.14	3.71 HI	(1.97, 6.34)	6.47
Lee, Leonard	449	192	1	0.52	1.47	0.63	(0.01, 3.52)	6.13
Lemaire, Anthony	167	122	6	4.92	1.84	4.78	(1.74, 10.39)	6.26
All Others (4)	501	231	2	0.87	1.20	1.29	(0.14, 4.66)	5.87
Anderson, Mark, B.**	107	20	0					
Ghaly, Aziz	48	33	0					
Prendergast, Thomas, W.**	160	98	0					
Scholz, Peter, M.**	186	80	2					

* = Operative Mortality includes: (1) all deaths occurring during the hospitalization in which the operation was performed, even after 30 days; and (2) those deaths occurring after discharge from the hospital, but within 30 days of the procedures.

** = Surgeon not currently performing CABG surgery in this hospital.

HI = Risk-adjusted mortality rate significantly higher than the New Jersey statewide mortality rate based on 95 percent confidence interval.

Table 2 (continued)

Risk-Adjusted Operative Mortality* Rate and Post-Surgery Length of Stay by Surgeon (2012-2013)

Hospital and Surgeon	Total Open Heart Procedures	Number of Isolated CABG Operations	Patient Operative Deaths*	Observed Patient Mortality(%)	Expected Patient Mortality(%)	Risk-Adjusted Patient Mortality (%)	95% Confidence Interval	Risk-Adjusted Post-Surgery Length of Stay
St Barnabas Medical Center								
Burns, Paul, G.**	326	148	2	1.35	1.49	1.62	(0.18, 5.87)	7.60
Sardari, Frederic, F.	366	208	3	1.44	1.60	1.61	(0.32, 4.71)	7.82
All Others (4)	33	12	0					
Camacho, Margarita	2	2	0					
Karanam, Ravindra, N.	3	2	0					
Russo, Mark	2	1	0					
Saunders, Craig, R.	26	7	0					
St Francis Medical Center								
All Others (9)	213	136	8	5.88	1.34	7.84 HI	(3.38, 15.45)	7.57
Anderson, Mark, B.**	46	4	0					
Batsides, George, P.**	2	2	0					
Lemaire, Anthony**	22	20	0					
Nixon, Todd**	5	4	0					
Prendergast, Thomas**	5	3	0					
Scholz, Peter, M.**	17	14	1					
Seinfeld, Fredric	30	26	0					
Shah, Salil**	13	10	3					
Shariff, Haji	73	53	4					
St Joseph's Hospital and Medical Center								
Asgarian, Kourosh	430	230	5	2.17	1.83	2.12	(0.68, 4.95)	4.47
Connolly, Mark, W.	287	166	3	1.81	2.02	1.60	(0.32, 4.67)	5.00
All Others (3)	169	79	2	2.53	1.51	2.99	(0.34, 10.81)	4.96
Badami, Chirag	27	20	0					
Christakos, Manny**	1	0	0					
Wohler, Alexander**	141	59	2					
St. Mary's Hospital (Passaic)								
All Others (2)	180	116	2	1.72	1.79	1.72	(0.19, 6.21)	6.28
Kaushik, Raj, R.**	133	78	1					
Shakir, Huzaifa, A.	47	38	1					
St Michael's Medical Center								
Patel, Nilesh, U.	261	159	2	1.26	1.20	1.88	(0.21, 6.78)	5.86
All Others (1)	43	19	0					
Krause, Tyrone, J.	43	19	0					
University Hospital								
All Others (2)	138	61	1	1.64	1.17	2.50	(0.03, 13.88)	6.25
Lovoulos, Constantinos	68	30	1					
Sambol, Justin, T.	70	31	0					
Valley Hospital								
Sperling, Jason, S.**	261	103	3	2.91	1.71	3.04	(0.61, 8.89)	5.21
Zapolanski, Alex	359	148	1	0.68	1.69	0.71	(0.01, 3.97)	5.07
All Others (2)	178	111	2	1.80	1.40	2.30	(0.26, 8.31)	5.15
Brizzio, Mariano	61	38	1					
Grau, Juan	117	73	1					
State Total (2012 - 2013)	15,848	7,616	136	1.79	1.79	1.79		6.19

* = Operative Mortality includes: (1) all deaths occurring during the hospitalization in which the operation was performed, even after 30 days; and (2) those deaths occurring after discharge from the hospital, but within 30 days of the procedures.

** = Surgeon not currently performing CABG surgery in this hospital.


HI = Risk-adjusted mortality rate significantly higher than the New Jersey statewide mortality rate based on 95 percent confidence interval.

Statewide Trends in Risk-adjusted Bypass Surgery Mortality Rates: Pooled Estimates

Figure 5 presents the statewide risk-adjusted mortality rates for years 1994 to 2013 derived by pooling data from all years (Sources: Appendix C; Appendix D, Table D3). When compared with 1994, the risk-adjusted operative mortality rate for bypass surgery in 2013 dropped 63.1 percent.

When a linear regression line is fitted to the pooled annual estimates, bypass mortality rate has been declining, in absolute terms, at the rate of 0.14 percentage points per year (See Appendix D, Figure D1).

Figure 5
Trends in Statewide Bypass Surgery Mortality Rates



Appendix A

Frequently Asked Questions

These are answers to some commonly asked questions that may be of interest to you as you read this report.

Q: Should I go only to the hospitals with below-average risk-adjusted mortality rates?

A: Not necessarily. There are many factors to consider in determining the best hospital for you. Among these are your own personal risk factors and the experience certain hospitals have treating patients with those risk factors. Before making up your mind, you should discuss this report with the physician, usually a cardiologist, who refers you for cardiac surgery. The cardiologist's knowledge and expertise will be a valuable guide in making your decision. You should also keep in mind that the data in this guide is from 2013 and that a hospital's performance may have changed since then.

Q: Should I avoid any surgeon whose volume is low in this report?

A: No, not necessarily. First, there are lower volume surgeons with good patient outcomes. Second, there may be a good explanation for why a surgeon had a low volume that is unrelated to his/her experience. For example, the surgeon may have recently moved from another state, where he/she performed a high volume of these procedures. It is best to discuss your concerns with your referring doctor.

Q: Should I refuse to go to a hospital or a surgeon for heart surgery if that hospital or surgeon has a worse than average mortality record?

A: Important decisions in areas such as cardiac surgery should be made after considering all available information. The statistics in this report are a starting point for discussions with your doctor. But they do not tell the complete story. That is why it is critical to bring your concerns and questions to your doctor.

Q: Is it better to go to a hospital with a high volume of cases?

A: National studies have demonstrated that, in general, hospitals with higher volumes have better results. However, some hospitals with high volumes have relatively high mortality rates, while others with low volumes have lower mortality rates.

Notes on Data:

The data used in this study were reported by hospitals according to criteria established by the Department, with assistance from the clinical experts. Additionally, the Department has made a good faith effort to ensure that the data elements and definitions are consistent with those issued by the Society for Thoracic Surgeons (STS). The data were audited by an independent reviewer under contract to the Department.

Throughout the process of developing this report, the Department has taken steps to make sure that all hospitals were informed about data reporting and auditing requirements, as well as the statistical methods being used to risk-adjust the reported mortality data.

The Department considers it a vital function of hospitals to be able to collect and report complete, accurate medical information on patients. This function is critical not only to the success of the cardiac surgery report, but to the hospitals' own ongoing efforts to improve the quality of care for all patients. The Department and hospitals will continue working to improve data collection procedures so that this report contains the best possible information.

Appendix B

New Jersey's Cardiovascular Health Advisory Panel (CHAP) Members

Perry Weinstock, MD
Chairperson of the CHAP

Director, Cooper Heart Institute
Chief of Cardiology

Cooper Hospital University Medical Center, Camden, New Jersey

Mary T. Abed, MD, FACC

Chief, Division of Cardiology
Jersey City Medical Center
Jersey City, New Jersey

Reginald J. Blaber, MD, FACC

Vice President, Cardiovascular Service Line
Our Lady of Lourdes Medical Center
Camden, New Jersey

Marc Cohen, MD, FACC

Chief, Division of Cardiology
Newark Beth Israel Medical Center
Newark, New Jersey

Pat Delaney, RN

Independent Consultant
North Haledon, New Jersey

Charles Dennis, MD, MBA, FACC

Independent Clinical and Management Consultant
Moorestown, New Jersey

Barry C. Esrig, MD, FACS, FACC, FCCP

New York Presbyterian Hospital
New York, New York

Austin Kutscher, Jr., MD, FACC

Hunterdon Cardiovascular Associates
Flemington, New Jersey

Glenn Laub, MD

Chair, Department of Cardiothoracic Surgery
Drexel University College of Medicine
Hahnemann University Hospital
Philadelphia, Pennsylvania

Leonard Lee, MD

Chair, Department of Surgery
Robert Wood Johnson Medical School
Chief, Division of Cardiothoracic Surgery
Robert Wood Johnson University Hospital
New Brunswick, New Jersey

Richard M. Niebart, MD

Jersey Shore University Medical Center
Mid-Atlantic Surgical Associates
Neptune, New Jersey

Grant V. S. Parr, MD, FACS, FACC, FCCP

Physician-in-Chief,
Gagnon Cardiovascular Institute
Atlantic Health
Morristown, New Jersey

Joseph E. Parrillo, MD

Professor of Medicine
Rutgers New Jersey Medical School
Chairman, Heart and Vascular Hospital
Hackensack University Medical Center
Hackensack, New Jersey

Department of Health Cardiac Surgery Report Team

Colette Lamothe-Galette

Director
Population Health Division

Abate Mammo, PhD

Executive Director
Healthcare Quality and Informatics

Jianping Huang, PhD

Priya Fox
Paul SanFilippo

Juana Jackson

Appendix C

Statewide Observed In-hospital and Operative Mortality Rates:

Year of Operation	Mortality Rate	
	In-hospital	Operative Mortality *
1994-1995	3.75	4.14
1996-1997	3.37	3.75
1998	2.60	3.01
1999	2.89	3.31
2000	2.22	2.68
2001	2.01	2.51
2002	1.80	2.15
2003	1.91	2.33
2004	1.54	1.98
2005	1.83	2.10
2006	1.73	2.00
2007	1.66	2.00
2008	1.19	1.47
2009	1.00	1.31
2010	1.58	1.95
2011	1.13	1.35
2012	1.63	2.01
2013	1.13	1.57

* Operative mortality includes the following:

- all deaths occurring during the hospitalization in which the operation was performed, even after 30 days; and
- deaths occurring after discharge from hospital, but within 30 days of the procedures.

Appendix D

Summary of Methods Used in this Report

Background

Five states, including New Jersey, have issued reports on isolated Coronary Artery Bypass Graft (CABG or bypass) surgery outcomes for hospitals and surgeons. New York first published a bypass surgery report in 1990 presenting 1989 data, with the latest report released in September 2016 using 2011-2013 data. Starting with its 1990 data, Pennsylvania has published several cardiac surgery reports, with its latest report released in November 2013 using 2011-2012 data. California has also published several cardiac surgery reports, with the most recent data released in January 2016 using 2013 data. Massachusetts published its first report on bypass surgery in October 2004 using 2002 data and released its latest report on a fiscal year basis (October 1, 2012 to September 30, 2013) in October 2015. In 1997, New Jersey began reporting on patient mortality for bypass surgery hospitals and surgeons, using 1994 and 1995 data combined.

The experience from these states is that disclosures have contributed to hospital quality improvement initiatives and significant reductions in bypass surgery mortality rate.

Factors That Affect a Patient's Risk of Bypass Surgery Mortality

The observed patient bypass surgery mortality rate for a hospital or surgeon is estimated as the number of bypass surgery patients who died in the hospital during or after surgery, or patients who died after discharge but within 30 days post-surgery, divided by the total number of patients who underwent the bypass surgery.

Unfortunately, this observed patient mortality rate is not a complete measure of the quality of care provided by a hospital or a

surgeon, because it does not account for how sick the patients were before surgery. If one hospital had considerably sicker patients than another hospital, it would be expected that its observed mortality rate would be somewhat higher. So it would not be fair to evaluate surgeons and hospitals performing bypass surgery solely on the basis of the percentage of their patients that died. For instance, an 85-year-old who had certain type of cerebrovascular disease and was in cardiogenic shock at the time of surgery would be at higher risk during this surgery than a 50-year-old who had no history of chronic disease.

To perform an even-handed analysis of the quality of surgical care provided by surgeons and hospitals performing bypass surgery, the Department adjusts the patient mortality rates for each surgeon and each hospital by the pre-surgery risk factors of each patient. This method gives hospitals and surgeons who operate on less healthy patients “extra credit.” Such hospitals and surgeons are not at a disadvantage when the outcome of the surgical care they provide is presented next to that of other hospitals and surgeons. Additionally, as stated earlier, extremely high risk patients, where the probability of death is very high, may, with the concurrence of the expert clinical panel, be excluded from the calculation.

The risk adjustment method is a statistical approach that uses results of a logistic regression analysis to assess the average risk of a bypass surgery for a patient. Key elements of the health histories of patients who have undergone bypass surgery in the same period, as well as their socio-demographic characteristics, are taken into account to estimate the expected outcome of a bypass surgery.

Assessing Patient Risk Factors

A logistic regression model which included all the before-surgery health and demographic factors was fitted to the data for the period covered by this report to identify those risk factors that were important in predicting whether a patient would die after a bypass surgery. The general form of a logistic regression model for estimating the “logit” of the probability of dying (p), denoted by Y_i , is presented as follows:

$$Y_i = \sum_k^K \beta_k X_{ki} + \varepsilon_i, \text{ Where } e^{X_{0i}} = 1;$$

$$Y_i = \log_e \left(\frac{p_i}{1 - p_i} \right) = \text{the "logit" of } p_i$$

- $i = 1, 2, \dots, n$; $k = 0, 1, 2, \dots, K$,
- β_k = Logistic regression coefficient for risk factor X_k ,
- K = Number of risk factors in the model,
- n = Number of patients,
- ε_i = Random error term i .

The statistically significant risk factors for this report (X_k) identified by the stepwise logistic regression analysis method are presented in Table D1. Table D1 also includes estimates of coefficients for the statistically significant risk factors, an indication of the level of statistical significance (p-values), and odds ratios. The list of risk factors includes only those that were statistically significant in predicting bypass surgery mortality with p-values of 0.05 or smaller.

The odds ratios are derived from the coefficients, and are used to compare the relative importance of the risk factors in predicting mortality from bypass surgery. For each of the risk factors identified in Table D1, the odds ratio represents how much as likely a patient is to die when compared to a patient who is in the reference group. So, for example, Table D1 shows that a patient who had cerebrovascular disease documented by non-invasive carotid test with > 79% occlusion before the surgery is more than two and half times (odds ratio = 3.53) as likely to die during or after bypass surgery compared to a patient who did not have the risk factor. This is based on the assumption that both patients have

the same set of other risk factors presented in the table.

Similarly, the odds of dying during or after bypass surgery for a patient who is in cardiogenic shock at the time of surgery is seven and three quarter times as likely (odds ratio= 7.75) compared with the odds of a patient who is not in cardiogenic shock at the time of surgery.

Estimation of Risk-adjusted Mortality Rates

The risk factors presented in Table D1 were used in the fitted logistic regression model to predict the probability of death from bypass surgery for each patient. The sum of predicted probabilities of dying for patients operated on in each hospital divided by the number of patients operated on in that hospital provides the predicted (or expected) death rate associated with the hospital. A similar analysis for a surgeon results in the expected death rate associated with that surgeon. Terms such as “expected” and “predicted” are used interchangeably in this report to signify that the estimates are derived from predicted probabilities after accounting for risk factors.

The predicted probability of dying for patient i (\hat{p}_i) is given as follows:

$$\hat{p}_i = \frac{e^{\hat{Y}_i}}{1 + e^{\hat{Y}_i}}, \text{ Where } i = 1, 2, 3, \dots, n; \text{ and}$$

$$\hat{Y}_i = \hat{\beta}_0 + \hat{\beta}_1 X_{1i} + \hat{\beta}_2 X_{2i} + \hat{\beta}_3 X_{3i} + \dots + \hat{\beta}_k X_{ki}$$

To assess the performance of each hospital or surgeon, we compared the observed patient mortality with the expected or predicted patient mortality, based on the existing risk factors for the hospital’s or surgeon’s patients. First, the observed patient mortality is divided by the expected mortality. If the resulting ratio is higher than one, the hospital or surgeon has a higher patient mortality than expected on the basis of their patient mix. If the ratio is lower than one, the hospital or surgeon has a lower mortality than expected, based on their patient mix. The ratio is then multiplied by the statewide mortality rate to produce the risk-

Cardiac Surgery in New Jersey

adjusted patient mortality rate for the hospital or the surgeon.

The risk-adjusted mortality rate represents the best estimate the fitted model provides using the statistically significant health risk factors. The risk-adjusted patient mortality rate represents what a hospital's or surgeon's patient mortality rate would have been if they had a mix of patients identical to the statewide mix. Thus, the risk-adjusted patient mortality has, to the extent possible, ironed out differences among hospitals and surgeons in patient mortality arising from the severity of illness of their patients.

The statistical methods described above are tested to determine if they are sufficiently accurate in predicting the risk of death for all patients – for those who are severely ill prior to

undergoing bypass surgery as well as those who are relatively healthy. In the analysis of data for this report, the tests confirmed that the model is reasonably accurate in predicting how patients of different risk levels will fare when undergoing bypass surgery. The area under the Receiver Operating Characteristic (ROC) curve, denoted by C-statistic in Table D1, was used to evaluate model performance. The C-statistic may be interpreted as the degree to which the risk factors in the model predicted the probability of death for bypass surgery patients. Specifically, the C-statistic measures the tendency of the predicted mortality for patients in the sample that died to be higher than that for patients who were discharged alive and were also alive 30 days after bypass surgery. The 2013 model C-statistic is 77.7 percent and is fairly high, suggesting that the model has strong predictive power.

Table D1
Risk Factors Identified for Isolated CABG Surgery Operative Mortality* (2013)

Patient Risk Factors Identified	Proportion of patients (%)	Logistic Regression Results		
		Coefficient	P-Value	Odds Ratio
Demographic factors				
Age	---	0.0378	0.0055	1.039
Health factors				
CVD - Non-Invasive>79%	1.24	1.2617	0.0292	3.531
Immunosuppressive Therapy	3.71	0.9255	0.0427	2.523
Peripheral Vascular Disease	13.68	0.9590	0.0009	2.609
Factors related to functioning of the heart				
Cardiogenic Shock	1.39	2.0482	<.0001	7.754
Congestive Heart Failure	21.33	0.8011	0.0037	2.228
NYHA Classification - IV	8.37	1.0243	0.0019	2.785
Intercept	-7.6226			
C-Statistic	0.777			
Number of CABGs (N)	3,881			

SOURCE: New Jersey Department of Health

* Operative Mortality includes: (1) all deaths occurring during the hospitalization in which the operation was performed, even after 30 days; and (2) those deaths occurring after discharge from the hospital, but within 30 days of the procedures.

Risk-adjusted Patient Mortality Rate Estimates

This section presents the results of our analysis including:

- (1) comparisons of risk-adjusted patient mortality rates for hospitals to the statewide rate in 2013;
- (2) comparisons of the statewide risk-adjusted patient mortality rate for each year in 1994-2013 to the rate for the whole period.

The risk-adjusted mortality rate estimates are presented in percentage points. The results also include the lowest and the highest risk-adjusted mortality rate estimates one would expect, using a 95 percent confidence level*.

* 95% confidence limits are calculated as follows:

$$LCL = \frac{D \left(1 - \frac{1}{9D} - \frac{1.96}{3\sqrt{D}} \right)^3}{E} S$$

$$UCL = \frac{(D+1) \left(1 - \frac{1}{9(D+1)} + \frac{1.96}{3\sqrt{(D+1)}} \right)^3}{E} S$$

Where *D* = Observed mortality, and
E = Predicted or Expected mortality,
S = Statewide rate.

(Source: Liddell, F. D. K., *Simple Exact Analysis of the Standardised Mortality Ratio. Journal of Epidemiology and Community Health*, 1984, 38, 85-88.)

Patient bypass surgery mortality rate by hospital compared with the statewide rate in 2013

The risk-adjusted patient mortality estimates from bypass surgery for each hospital in 2013 are presented in Table D2. The results compare each hospital's risk-adjusted patient mortality rate with the statewide mortality rate.

After adjusting for how sick the patients were before surgery at each hospital, we present the estimates of risk-adjusted patient mortality rate for each hospital in the sixth column of Table D2.

If a hospital's 95 percent confidence interval contains the statewide rate, it means that the difference between the hospital's risk-adjusted mortality rate and the statewide rate was not statistically significant. If the whole of a hospital's 95 percent confidence interval clearly falls to the left of the statewide rate, it means that the hospital's risk-adjusted patient mortality rate was statistically significantly lower than the statewide rate. If the whole of the 95 percent confidence interval falls to the right of the statewide rate, it means that the hospital's risk-adjusted mortality rate was statistically significantly higher than the statewide rate.

The observed operative mortality rate statewide in 2013 for bypass patients was 1.57 percent, based on 61 deaths out of 3,881 bypass operations performed. Table D2 presents the bypass volume, observed mortality rate, expected mortality rate, risk-adjusted mortality rate and its confidence interval, as well as risk-adjusted length of stay following bypass surgery for each of the 18 hospitals.

In 2013, one hospital, St. Francis Medical Center, had a statistically significantly higher risk-adjusted mortality rate than the statewide rate.

Table D2

Patient Operative Mortality* and Length of Stay After Isolated CABG Surgery by Hospital (2013)

Hospital	Number of Isolated CABG Operations	Patient Operative Deaths*	Observed Patient Mortality (%)	Expected Patient Mortality (%)	Risk-Adjusted Patient Mortality (%)	95% Confidence Interval	Risk-Adjusted Patient Post-Surgery LOS (days)
AtlantiCare Regional Medical Center	148	1	0.68	1.58	0.67	(0.01, 3.74)	5.77
Cooper Hospital University Medical Center	172	5	2.91	1.82	2.51	(0.81, 5.86)	5.25
Deborah Heart and Lung Center	119	2	1.68	2.09	1.26	(0.14, 4.56)	5.87
Englewood Hospital and Medical Center	157	0	0.00	1.66	0.00	(0.00, 2.21)	6.33
Hackensack University Medical Center	278	1	0.36	1.68	0.34	(0.00, 1.88)	7.60
Jersey City Medical Center	147	3	2.04	1.34	2.39	(0.48, 6.97)	5.34
Jersey Shore University Medical Center	394	8	2.03	1.59	2.00	(0.86, 3.95)	6.72
Morristown Medical Center	643	6	0.93	1.58	0.93	(0.34, 2.02)	6.33
Newark Beth Israel Medical Center	168	3	1.79	1.62	1.73	(0.35, 5.06)	6.95
Our Lady of Lourdes Medical Center	389	7	1.80	2.22	1.27	(0.51, 2.62)	5.48
Robert Wood Johnson University Hospital	420	8	1.90	1.13	2.66	(1.15, 5.24)	6.26
Saint Barnabas Medical Center	182	1	0.55	1.26	0.69	(0.01, 3.81)	7.59
St. Francis Medical Center	75	7	9.33	1.04	14.08	HI (5.64, 29.02)	7.04
St. Joseph's Hospital and Medical Center	234	3	1.28	1.60	1.26	(0.25, 3.69)	4.56
St. Mary's Hospital (Passaic)	64	1	1.56	1.29	1.91	(0.02, 10.63)	6.35
St. Michael's Medical Center	79	1	1.27	0.72	2.76	(0.04, 15.36)	5.91
University Hospital	29	0	0.00	0.84	0.00	(0.00, 23.61)	6.60
Valley Hospital	183	4	2.19	1.54	2.23	(0.60, 5.70)	5.10
Statewide	3,881	61	1.57	1.57	1.57		6.12

SOURCE: New Jersey Department of Health

* = Operative Mortality includes: (1) all deaths occurring during the hospitalization in which the operation was performed, even after 30 days; and (2) those deaths occurring after discharge from the hospital, but within 30 days of the procedures.

HI - Risk-adjusted mortality rate significantly higher than New Jersey statewide mortality rate based on 95 percent confidence interval.

Annual risk-adjusted mortality compared to the combined 1994-2013 risk-adjusted mortality

Table D3 presents the results of an analysis to identify the trend in the statewide mortality rate of patients who underwent bypass surgery using a statistical model based on the pooled data collected over the period 1994-2013. For each of the years, the table presents the observed patient mortality rate, the expected patient mortality rate, and the statewide risk-adjusted patient mortality rate estimate. Note that the numbers differ from those shown in reports produced in previous years, due to the revised definition of mortality and the use of pooled data for the analysis. The table further exhibits whether the risk-adjusted mortality rate for the year is statistically different from the pooled mortality rate for the 1994-2013 period.

Table D3 also shows that between 2012 and 2013, the number of bypass surgeries performed in New Jersey increased from 3,735 to 3,881 or by 3.9 percent. Over the same time period, the number of deaths decreased from 75 to 61 or by 18.7 percent. On a risk-adjusted basis, the mortality rate decreased 16.5 percent between 2012 and 2013, which was not statistically significant. Nevertheless, since 1994 risk-adjusted mortality rate has declined 63.1 percent, which is statistically significant.

The trend in operative bypass mortality rate between 1994 and 2013 was estimated by fitting a regression line of pooled annual risk-adjusted bypass mortality rates to procedure year (Figure D1). According to the fitted regression line, operative mortality from bypass surgery has been declining, in absolute terms, at the rate of 0.14 percentage points per year between 1994 and 2013 ($R^2 = 0.81$).

Table D3
Annual Risk-Adjusted Patient Operative Mortality Rate* Derived from the Pooled Data for the Period 1994-2013

Year	Number of Isolated CABG Operations	Operative Patient Mortality*	Observed Patient Mortality Rate (%)	Predicted Patient Mortality Rate (%)	Risk-Adjusted Patient Mortality Rate (%)		Yearly Change in Risk-Adjusted Mortality Rate (%)	Percent Change from 1994 Risk-Adjusted Mortality Rate (%)
1994	6,957	274	3.94	2.40	4.37	HI	----	----
1995	7,553	327	4.33	2.59	4.45	HI	0.08	1.8
1996	8,262	341	4.13	2.74	4.01	HI	-0.44	-8.2
1997	8,286	280	3.38	2.87	3.14	HI	-0.87	-28.2
1998	8,377	252	3.01	2.78	2.88	SA	-0.26	-34.1
1999	8,108	268	3.31	2.95	2.99	SA	0.10	-31.7
2000	8,220	220	2.68	2.62	2.72	SA	-0.27	-37.8
2001	8,045	202	2.51	2.69	2.49	SA	-0.23	-43.0
2002	7,391	159	2.15	2.70	2.12	LO	-0.37	-51.5
2003	6,817	159	2.33	2.71	2.29	SA	0.17	-47.6
2004	6,177	122	1.98	2.57	2.04	LO	-0.25	-53.2
2005	5,576	117	2.10	2.56	2.18	LO	0.14	-50.1
2006	5,211	104	2.00	2.56	2.08	LO	-0.10	-52.4
2007	4,943	99	2.00	2.59	2.06	LO	-0.02	-52.9
2008	4,620	68	1.47	2.53	1.55	LO	-0.51	-64.5
2009	4,497	59	1.31	2.49	1.40	LO	-0.15	-67.9
2010	4,302	84	1.95	2.70	1.92	LO	0.52	-56.0
2011	3,709	50	1.35	2.52	1.43	LO	-0.50	-67.3
2012	3,735	75	2.01	2.77	1.93	LO	0.50	-55.8
2013	3,881	61	1.57	2.60	1.61	LO	-0.32	-63.1
1994-2013	124,667	3,321	2.66	2.66	2.66			

SOURCE: New Jersey Department of Health


* Operative Mortality includes: (1) all deaths occurring during the hospitalization in which the operation was performed, even after 30 days; and (2) those deaths occurring after discharge from the hospital, but within 30 days of the procedures.

LO - The risk-adjusted patient mortality is significantly lower than the mortality for the 1994-2013 combined when evaluated with a 95 percent confidence interval.

SA - The risk-adjusted patient mortality is same as the mortality for the 1994-2013 combined when evaluated with a 95 percent confidence interval.

HI - The risk-adjusted patient mortality is significantly higher than the mortality for the 1994-2013 combined when evaluated with a 95 percent confidence interval.

Figure D1
Trend in Risk-Adjusted Operative Mortality* Rate (1994-2013)



SOURCE: New Jersey Department of Health

* Operative Mortality includes: (1) all deaths occurring during the hospitalization in which the operation was performed, even after 30 days; and (2) those deaths occurring after discharge from the hospital, but within 30 days of the procedures.

Risk factors for post-surgery length of stay

In an attempt to predict a patient’s post-operative length of stay, we fitted a generalized linear regression model on the log transformation of length of stay. The model was developed using demographic factors, health factors, factors related to functioning of the heart and prior cardiac intervention as predictors. Patients who died during the bypass surgery hospitalization were excluded from analysis as were patients who stayed fewer than two days in hospital and those who stayed over 30 days.

Table D4 presents the final model used to estimate risk-adjusted length of stay by hospital and includes only those predictors found to be statistically significant at five percent or lower levels. Consistent with findings in Pennsylvania, the predictive power of the model is low (only 14.0 percent). Such low predictive power is usually common when one fits a regression model using individual level data as large as these.

Please note that the coefficients provided in Table D4 are in log form and interpretation of the values should take that into consideration.

Table D4
Risk Factors Identified for Isolated CABG Surgery Length of Stay (2013)

Patient Risk Factors Identified	Proportion of Patients(%)	Generalized Linear Regression Results	
		Coefficient	P-Value
Demographic factors			
Ages 55 to 59	12.43	0.05949	0.0134
Ages 60 to 64	16.11	0.07108	0.0017
Ages 65 to 69	18.64	0.09997	<.0001
Ages 70 to 74	15.06	0.15172	<.0001
Ages 75 to 79	11.70	0.17940	<.0001
Ages 80 to 84	8.39	0.25272	<.0001
Ages 85 and over	3.15	0.34631	<.0001
Female	24.82	0.05185	0.0005
Non-hispanic Other	9.20	0.05809	0.0074
Health factors			
Diabetes - Insulin	15.64	0.07466	<.0001
Immunosuppressive Therapy	3.60	0.07706	0.0226
Lung Disease	18.96	0.04704	0.0037
Obesity	13.07	0.08745	<.0001
Renal Failure without Dialysis	3.00	0.14649	<.0001
Renal Failure with Dialysis	2.73	0.14439	0.0002
Factors related to functioning of the heart			
Angina - Unstable	43.27	0.06318	<.0001
Arrhythmia - Heart Block	0.66	0.15975	0.0383
Arrhythmia - AFib/Flutter	7.60	0.08276	0.0005
Cardiogenic Shock - Hemodynamic Unstability	1.03	0.18362	0.0049
Congestive Heart Failure	20.85	0.09689	<.0001
Ejection Fraction 1 - 29%	5.78	0.10565	0.0002
Ejection Fraction 30 - 49%	26.37	0.04259	0.0037
	34.75	0.02697	0.0448
Number of Diseased Vessels - Two	19.58	0.10137	0.0030
Number of Diseased Vessels - Three	76.29	0.15933	<.0001
PTCA Received <= 6 Hours	0.79	0.14809	0.0395
Resuscitation	0.26	0.38640	0.0018
IABP Preop	7.83	0.08027	0.0015
Intercept	1.4035		
R-Square	14.04		
Number of CABGs (N)*	3,804		

SOURCE: New Jersey Department of Health

* Excluded are patients who died during hospitalization where CABG was performed; patients with postsurgical LOS > 30 days; and patients with post-surgical LOS < 2 days.

References

- The Massachusetts Department of Public Health, Adult Coronary Artery Bypass Graft Surgery in the Commonwealth of Massachusetts, Fiscal Year 2013 Report (October 1, 2012 – September 30, 2013). Massachusetts Data Analysis Center, Department of Health Care Policy, Harvard Medical School, October 2015.
<http://www.massdac.org/wp-content/uploads/CABG-FY2013.pdf>
- New Jersey Department of Health, Cardiac Surgery in New Jersey, Health Care Quality Assessment, Office of Policy and Strategic Planning, December 2015.
<http://www.nj.gov/health/healthcarequality/documents/cardconsumer15.pdf>
- New York State Department of Health, New York State Department of Health, Adult Cardiac Surgery in New York State: 2011-2013, September 2016.
http://www.health.ny.gov/statistics/diseases/cardiovascular/heart_disease/docs/2011-2013_adult_cardiac_surgery.pdf
- California Office of Statewide Health Planning and Development, Coronary Artery Bypass Graft Surgery: 2013 Data. Sacramento, CA, January 2016.
http://www.oshpd.ca.gov/HID/Products/Clinical_Data/CABG/13Breakdown.html
- Pennsylvania Health Care Cost Containment Council, Cardiac Surgery in Pennsylvania, Information about Hospitals and Cardiothoracic Surgeons, Data: July 1, 2011 to December 31, 2012: Information about hospitals and cardiothoracic surgeons. November 2013.
<http://www.phc4.org/reports/cabg/12/docs/cabg2012report.pdf>

Limited copies are available by writing to the New Jersey Department of Health, Office of Health Care Quality Assessment, P.O. Box 360, Trenton, NJ 08625; or by phone at (800) 418-1397; or fax at (609) 984-7735; or email to hcqa@doh.state.nj.us. The report is also posted on our website at www.nj.gov/health/healthcarequality/cardiacsurgery.shtml.



**Cardiac
Surgery in
New Jersey, 2013**

