

## SUMMARY OF RESULTS – FIBI002



1. Stream Name:	Furnace Bk
2. Sampling Date:	9/19/00
3. Sampling Location:	Pequest WMA @ Old Railroad Crossing (40 49 38.71N; 74 58 50.00W)
4. County:	Warren
5. Watershed Management Area:	1
6. Contributing Drainage Area (Sq. Mi.):	7.3
7. Stream Water Quality Class:	FW2-NT
8. FIBI Rating:	Good (40) (See Appendix 3)
9. Habitat Assessment Rating:	Optimal (189) (See Appendix 3)
10. Fishable Species Present:	Yes
11. Relevant AMNET <sup>1</sup> Station Data:	
Proximity of FIBI station to AMNET station:	.6mi downstream AN0042
AMNET Rating:	1992- Severe; 1997 - Moderate
12. Stream Chemistries:	
Dissolved Oxygen (mg/l)	9.48
Temperature °C.	15.4
pH	8.62
Conductivity (µmhos/cm)	596
Secchi Disk (inches)	NA
13. Number of Fish With Anomalies:	0
14. Water Clarity:	clear
15. Forest Canopy:	Partially Open
16. Flow:	moderate
17. Substrate: (qualitative)	20% Gravel/Sand, 20% Cobble, 50% Boulder, 10% Silt
18. Habitat Type: (qualitative)	30% Riffle, 60% Run, 10% Pool
19. Other observations:	
20. Number of Fish Species Identified: (see next page)	15
21. Total Number of Fish Collected:	360

<sup>1</sup> AMNET is the acronym for the DEP's ambient benthic macroinvertebrate monitoring network – a series of 820 monitoring stations located throughout the state's waterways that collects data on the health of bottom dwelling stream fauna which in turn is used to assess general water quality.

LISTED IN ORDER OF ABUNDANCE FOUND (see also Figure 1.1)

COMMON NAME	SCIENTIFIC NAME	# FOUND	SIZE RANGE (INCHES)
White Sucker*	<i>Catostomus commersoni</i>	88	
Blacknose Dace	<i>Rhinichthys atratulus</i>	75	
Creek Chub	<i>Semotilus atromaculatus</i>	66	
Tessellated Darter	<i>Etheostoma olmstedii</i>	43	
Bluegill*	<i>Lepomis macrochirus</i>	38	1.6 - 4.7
American Eel*	<i>Anguilla rostrata</i>	20	
Sea Lamprey	<i>Petromyzon marinus</i>	7	
Longnose Dace	<i>Rhinichthys cataractae</i>	7	
Redfin Pickerel*	<i>Esox americanus americanus</i>	3	3.9 - 5.5
Redbreast Sunfish*	<i>Lepomis auritus</i>	3	1.6 - 5.5
Banded Killifish	<i>Fundulus diaphanus</i>	3	
Yellow Perch*	<i>Perca flavescens</i>	2	4.3 - 4.7
Pumpkinseed*	<i>Lepomis gibbosus</i>	2	3.1 - 3.1
Largemouth Bass*	<i>Micropterus salmoides</i>	2	3.1 - 3.9
Spotfin Shiner	<i>Cyprinella spiloptera</i>	1	

\*Regulated as a fishable species under current New Jersey Fish and Wildlife codes

**HABITAT ASSESSMENT FOR HIGH GRADIENT STREAMS FURNACE BK (FIBI002) – 9/19/00**

Habitat Parameter	Condition Category			
	Optimal	Suboptimal	Marginal	Poor
<b>1. Epifaunal Substrate/Available Cover</b> Greater than 50% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are <b>not</b> new fall and not transient). SCORE 19	20 <b>19</b> 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
<b>2. Embeddedness</b> Gravel, cobble, and boulder particles are 0-25% surrounded by fine sediment. SCORE 18	20 19 <b>18</b> 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
<b>3. Riffle Quality</b> Well-developed riffle and run; riffle is as wide as stream and length extends two times the width of stream; abundance of cobble. (Boulders prevalent in headwater streams). SCORE 17	20 19 18 <b>17</b> 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
<b>4. Sediment Deposition</b> Little or no enlargement of islands or point bars and less than 5% (<20% for low-gradient streams) of the bottom affected by sediment deposition. SCORE 20	<b>20</b> 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
<b>5. Channel Flow Status</b> Water reaches base of both lower banks, and minimal amount of channel substrate is exposed. SCORE 19	20 <b>19</b> 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
<b>6. Channel Alteration</b> Channelization or dredging absent or minimal; stream with normal pattern. SCORE 20	<b>20</b> 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
<b>7. Frequency of Riffles (or bends)</b> Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important. All 4 velocity/depth patterns present. SCORE 18	20 19 <b>18</b> 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
<b>8. Bank Stability (score each bank)</b> Note: determine left or right side by facing downstream. SCORE <u>  </u> 9 (LB) SCORE 9 (RB)	Left Bank 10 <b>9</b> Right Bank 10 <b>9</b>	8 7 6 8 7 6	5 4 3 5 4 3	2 1 0 2 1 0
<b>9. Bank Vegetative Protection (score each bank)</b> More than 90% of the streambank surfaces and immediate riparian zone covered by native vegetation, including trees, understory shrubs, or nonwoody macrophytes; vegetative disruption through grazing or mowing minimal or not evident; almost all plants allowed to grow naturally. SCORE 10 (LB) SCORE 10 (RB)	Left Bank <b>10</b> 9 Right Bank <b>10</b> 9	8 7 6 8 7 6	5 4 3 5 4 3	2 1 0 2 1 0
<b>10. Riparian Vegetative Zone Width (score each bank riparian zone)</b> Width of riparian zone >18 meters; human activities (i.e., parking lots, roadbeds, clear-cuts, lawns, or crops) have not impacted zone. SCORE 10 (LB) SCORE 10 (RB)	Left Bank <b>10</b> 9 Right Bank <b>10</b> 9	8 7 6 8 7 6	5 4 3 5 4 3	2 1 0 2 1 0

**HABITAT SCORE**  
  
 189

HABITAT SCORES	VALUE
OPTIMAL	160 C 200
SUB-OPTIMAL	110 C 159
MARGINAL	60 C 109
POOR	< 60

**FIBI002 - Furnace Bk @ Pequest Rd**  
**Date Sampled - 9/19/2000**

Excellent **Good** Fair Poor

	<b>Score</b>
# of Fish Species	5
# of Benthic Insectivorous Species (BI)	5
# of Trout and Centrarchid Species (trout, bass, sunfish, crappie)	5
# of Intolerant Species (IS)	1
Proportion of Individuals as White Suckers	3
Proportion of Individuals as Generalists (carp, creek chub, banded killifish, goldfish, fathead minnow, green sunfish)	5
Proportion of Individuals as Insectivorous <b>Cyprinids</b> (I and BI)	3
Proportion of Individuals as Trout	
OR	
Proportion of Individuals as Piscivores (Excluding American Eel)*	3
Number of Individuals in Sample	5
Proportion of Individuals w/disease/anomalies (excluding blackspot)	5
<b>Total</b>	<b>40</b>

**Stream Rating**

<b>45-50</b>	<b>Excellent</b>
<b>37-44</b>	<b>Good</b>
<b>29-36</b>	<b>Fair</b>
<b>10-28</b>	<b>Poor</b>