



Pohatcong Creek - FIBI033

Drainage Area of FIBI033: 9.8 Square Miles

Surface Water Quality Classification of FIBI033: FW2-TM



	FIBI Sampling Location
	Small Streams (1st and 2nd Order)
	Large Streams (3rd Order and Above)



SUMMARY OF RESULTS – FIBI033



1. Stream Name:	Pohatcong Creek
2. Sampling Date:	07/31/2001
3. Sampling Location:	SR 31 (40 46 52N; 74 58 29W)
4. Municipality	Washington Twp.
5. County:	Warren
6. Watershed Management Area:	1
7. Contributing Drainage Area (Sq. Mi.):	9.8
8. Stream Water Quality Class:	FW2-TM
9. FIBI Rating:	Good (44) (See Appendix 3)
10. Habitat Assessment Rating:	Suboptimal (145) (See Appendix 3)
11. Fishable Species Present:	Yes
12. Relevant AMNET ¹ Station Data:	
Proximity of FIBI station to AMNET station:	0.94 mi. downstream of AN0055
AMNET Rating:	1992-Moderately Impaired; 1997-Moderately Impaired
13. Stream Chemistries:	
Dissolved Oxygen (mg/l)	9.2
Temperature °C.	19.8
pH	8
Conductivity (µmhos/cm)	231
14. Number of Fish With Anomalies:	0
15. Length of Stream Segment Sampled	150 meters (492 feet)
16. Water Clarity:	Clear
17. Average Forest Open Canopy:	28%
18. Discharge (ft. ³ /sec.):	23.7
19. Substrate: (qualitative)	5% Gravel/Sand, 60% Cobble, 30% Boulder, 5% Silt
20. Habitat Type: (qualitative)	35% Riffle, 15% Run, 50% Pool
21. Other observations:	N/A
22. Number of Fish Species Identified: (see next page)	22
23. Total Number of Fish Collected:	667

¹ AMNET is the acronym for the DEP's ambient benthic macroinvertebrate monitoring network – a series of 820 monitoring stations located throughout the state's waterways that collects data on the health of bottom dwelling stream fauna which in turn is used to assess general water quality.

FIBI033
POHATCONG CREEK
SR 31
Washington Twp., Warren Co.

SR 31

Jackson Valley Rd



0 0.1 Miles



LEGEND	
#	Start
#	Finish
—	Segment Sampled
Ⓜ	Direction of Flow



FIBI033 - Pohatcong Creek @ Rt. 31
Date Sampled - 7/31/2001

Excellent **Good** Fair Poor

	Score
# of Fish Species	5
# of Benthic Insectivorous Species (BI)	5
# of Trout and Centrarchid Species (trout, bass, sunfish, crappie)	5
# of Intolerant Species (IS)	5
Proportion of Individuals as White Suckers	3
Proportion of Individuals as Generalists (carp, creek chub, banded killifish, goldfish, fathead minnow, green sunfish)	5
Proportion of Individuals as Insectivorous Cyprinids (I and BI)	5
Proportion of Individuals as Trout	
OR	
Proportion of Individuals as Piscivores (Excluding American Eel)*	1
Number of Individuals in Sample	5
Proportion of Individuals w/disease/anomalies (excluding blackspot)	5
Total	44

Stream Rating

45-50 Excellent
37-44 Good
29-36 Fair
10-28 Poor

HABITAT ASSESSMENT FOR HIGH GRADIENT STREAMS Pohatcong Creek (FIBI033) – 7/31/01

	Condition Category			
	Optimal	Suboptimal	Marginal	Poor
1. Epifaunal Substrate /Available Cover Greater than 70% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are <u>not</u> new fall and <u>not</u> transient). SCORE 13	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
2. Embeddedness Gravel, cobble, and boulder particles are 0-25% surrounded by fine sediment. Layering of cobble provides diversity of niche space SCORE 16	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
3. Velocity/Depth Regimes All 4 velocity/depth regimes present (slow-deep, slow-shallow, fast-deep, fast-shallow). (slow is <0.3 m/s, deep is >0.5 m) SCORE 17	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
4. Sediment Deposition Little or no enlargement of islands or point bars and less than 5% (<20% for low-gradient streams) of the bottom affected by sediment deposition. SCORE 13	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
5. Channel Flow Status Water reaches base of both lower banks, and minimal amount of channel substrate is exposed. SCORE 20	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
6. Channel Alteration Channelization or dredging absent or minimal; stream with normal pattern. SCORE 19	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
7. Frequency of Riffles (or bends) Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important. SCORE 12	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
8. Bank Stability (score each bank) Note: determine left or right side by facing downstream. SCORE <u>6</u> (LB) SCORE <u>7</u> (RB)	Left Bank 10 9 Right Bank 10 9	8 7 6 8 7 6	5 4 3 5 4 3	2 1 0 2 1 0
9. Bank Vegetative Protection (score each bank) SCORE <u>3</u> (LB) SCORE <u>9</u> (RB)	Left Bank 10 9 Right Bank 10 9	8 7 6 8 7 6	5 4 3 5 4 3	2 1 0 2 1 0
10. Riparian Vegetative Zone Width (score each bank riparian zone) SCORE <u>0</u> (LB) SCORE <u>10</u> (RB)	Left Bank 10 9 Right Bank 10 9	8 7 6 8 7 6	5 4 3 5 4 3	2 1 0 2 1 0

HABITAT SCORE

145

HABITAT SCORES	VALUE
OPTIMAL	160 C 200
SUB-OPTIMAL	110 C 159
MARGINAL	60 C 109
POOR	< 60

FIBI033 07/31/01

POHATCONG CREEK

LISTED IN ORDER OF ABUNDANCE FOUND

COMMON NAME	SCIENTIFIC NAME	# FOUND	SIZE RANGE (INCHES)
Blacknose Dace	<i>Rhinichthys atratulus</i>	138	
Common Shiner	<i>Luxilus cornutus</i>	133	
White Sucker*	<i>Catostomus commersoni</i>	112	
Cutlips Minnow	<i>Exoglossum maxillingua</i>	54	
Redbreast Sunfish*	<i>Lepomis auritus</i>	50	1.6 - 5.3
Satinfin Shiner	<i>Cyprinella analostana</i>	38	
Tesselated Darter	<i>Etheostoma olmstedii</i>	33	
Longnose Dace	<i>Rhinichthys cataractae</i>	18	
Spottail Shiner	<i>Notropis hudsonius</i>	15	
Sea Lamprey	<i>Petromyzon marinus</i>	13	
Fallfish	<i>Semotilus corporalis</i>	12	
Brown Trout*	<i>Salmo trutta</i>	12	2.0 - 11.8
American Eel*	<i>Anguilla rostrata</i>	11	
Rock Bass*	<i>Ambloplites rupestris</i>	10	3.0 - 6.7
Margined Madtom	<i>Noturus insignis</i>	7	
Creek Chub	<i>Semotilus atromaculatus</i>	3	
Bluegill*	<i>Lepomis macrochirus</i>	3	3.3
Yellow Bullhead*	<i>Ameiurus natalis</i>	1	4.7
Pumpkinseed*	<i>Lepomis gibbosus</i>	1	3.3
Creek Chubsucker	<i>Erimyzon oblongus</i>	1	
Rainbow Trout*	<i>Oncorhynchus mykiss</i>	1	9.8
Brook Trout*	<i>Salvelinus fontinalis</i>	1	8.3

* Regulated as a fishable species under current New Jersey Fish and Wildlife codes


**FIGURE 1.1 (Not To Scale)
Species Identified at Pohatcong Creek (FIBI033)**

John Scarola



White Sucker

John Scarola




Rainbow Trout

John Scarola




Brown Trout

John Scarola




Sea Lamprey

John Scarola



American Eel

John Scarola



Fallfish


**FIGURE 1.1 (Not To Scale)
Species Identified at Pohatcong Creek (FIBI033)**

John Scarola



Common Shiner

Jenkins & Burkhead



Cutlips Minnow

Konrad Schmidt



Spottail Shiner

John Scarola



Blacknose Dace

Jenkins & Burkhead



Satinfin Shiner

Shute



Longnose Dace

**FIGURE 1.1 (Not To Scale)
Species Identified at Pohatcong Creek (FIBI033)**

John Scarola



Rock Bass

Shute



Margined Madtom

Konrad Schmidt



Creek Chub

John Scarola



Redbreast Sunfish

William Roston



Creek Chubsucker

John Scarola



Pumpkinseed

**FIGURE 1.1 (Not To Scale)
Species Identified at Pohatcong Creek (FIBI033)**

Don Beimborn




Brook Trout

John Scarola



Yellow Bullhead

John Scarola



Tessellated Darter

John Scarola



Bluegill