

Patient Safety Indicators

Technical Report

A Supplement to the

Hospital Performance Report - 2012

Published in 2015

Health Care Quality Assessment

**Office of Policy and Strategic Planning
New Jersey Department of Health**

* For inquiries, contact Markos Ezra, PhD, by phone at (800) 418-1397 or by email at Markos.Ezra@doh.state.nj.us.

Table of Contents

	Page
Executive Summary	ii
Introduction	1
The Patient Safety Indicators (PSI) Module	2
How are PSI rates calculated?	3
Interpretation of PSI rates	4
Limitations of PSIs measures	6
Patient Safety Indicator estimates for New Jersey	7
Statewide PSI estimates compared to National estimates	41
Summary of findings.....	42
References.....	43

Tables

Table 1. Retained surgical item or unretrieved device fragment count.....	9
Table 2. Iatrogenic pneumothorax (per 1,000 per 1,000 med. & surg. discharges).....	11
Table 3. Postoperative hip fracture (per 1,000 surgical discharges)	14
Table 4. Perioperative hemorrhage or hematoma (per 1,000 surgical discharges)	17
Table 5. Perioperative pulmonary embolism or deep vein thrombosis (per 1,000 surgical discharges).....	20
Table 6. Postoperative sepsis (per 1,000 elective surgical discharges)	23
Table 7. Postoperative wound dehiscence (per 1,000 abdominopelvic surgical discharges)	26
Table 8. Accidental puncture or laceration (per 1,000 med. & surg. discharges)	29
Table 9. Birth trauma - injury to neonate (per 1,000 live births)	32
Table 10. Obstetric trauma - vaginal with instrument (per 1,000 instrument-assisted vaginal deliveries)	35
Table 11. Obstetric trauma - vaginal w/o instrument (per 1,000 vaginal deliveries).....	38
Table 12. Comparing New Jersey's Statewide PSI Rates with National Rates	41

Executive Summary

The Office of Health Care Quality Assessment (HCQA) of the New Jersey Department of Health and Senior Services assesses health care quality using qualitative and quantitative data reported by hospitals to support performance monitoring related to patient care and safety. Specifically, HCQA produces consumer reports on cardiac surgery, hospital performance, and hospital quality indicators; reviews confidential reports and root-cause analyses of reportable medical errors; and maintains several databases to support licensure requirements. In an effort to enhance the information the Department provides to the public on hospital care, HCQA staff apply statistical tools developed by the Federal Agency for Healthcare Research and Quality (AHRQ) to the New Jersey hospital discharge data commonly known as Uniform Billing (UB) data. This report, presents findings resulting from the application of a statistical tool known as the Patient Safety Indicator (PSI) module to the 2012 New Jersey hospital discharge data. The PSI module currently in use contains 18 hospital-level and seven area-level indicators that reflect the quality of care by hospitals. These indicators serve as flags for potential quality problems (adverse events) rather than provide definitive measures of quality of care. According to the AHRQ, the 18 indicators are selected based on their ability to screen out conditions present on admission from conditions that develop after admission, the potential preventability of the complication, and the ability of the indicator to identify medical error.

This report is a supplement to the Hospital Performance Report and covers only the 12 PSIs mandated for public reporting by law. In 2009, legislation (S2471) was signed into law requiring that hospital-specific data on patient-safety performance and serious medical errors be included in the annual New Jersey Hospital Performance Report. The data in this report present adverse events during hospitalization in each of the 72 licensed hospitals currently operating in the state. For the seven PSIs, risk-adjusted rates are provided along with confidence intervals to help make a statistical assessment of patient safety in the hospital. Statewide and national estimates are also provided to help compare hospital performance to the state or to the national rates.

Comparison of a hospital's rate to the statewide rate (presented in the top row of each of the PSI tables (Tables 1-11) is one way to assess how well that hospital performed among its peers in the state. A hospital's peers could be defined at many levels (e.g., teaching hospitals, urban hospitals, suburban hospitals, etc.). It is suggested that a hospital's performance be assessed by looking at its performance across the several PSI estimates presented in the PSI tables.

The 2012 New Jersey data show that there are substantial variations in risk-adjusted rates of adverse events by hospital. Some hospitals exhibit significantly higher risk-adjusted rates (risk-adjusted adverse event rates) than the

corresponding statewide rates while others have significantly lower rates than the statewide rates.

Some Highlights

- Statewide, in 2012, there were a total of 38 cases identified as '*Retained surgical item or unretrieved device fragment*' (formerly *foreign body left during procedure*). These 38 events were reported by 21 hospitals, with three hospitals (Hackensack University MC, RWJ University Hospital and UMDNJ-University Hospital) reporting 17 of the 38 cases. Table 1 shows the distribution of these adverse events by hospital.
- Overall, there were 238 cases of '*iatrogenic pneumothorax*' in 2012, for a risk-adjusted rate of 0.3 per 1,000 medical and surgical discharges. Hospital-specific rates for this indicator ranged from a low of 0.0 to a high of 1.4 per 1,000 medical and surgical discharges.
- Statewide, there were 1,086 '*Perioperative hemorrhage or hematoma*' cases in 2012, for a risk-adjusted rate of 14.7 per 1,000 surgical discharges. Hospital-specific rates for this indicator ranged from a low of 0.0 to a high of 28.5 per 1,000 surgical discharges.
- Statewide, there were 1,315 cases of '*Perioperative deep vein thrombosis (DVT) or pulmonary embolism (PE)*' in 2012, for a risk-adjusted rate of 5.5 per 1,000 surgical discharges. Hospital-specific rates for this indicator ranged from a low of 0.0 to a high of 42.9 per 1,000 surgical discharges (see Table 5).
- Statewide, there were 157 '*Postoperative sepsis*' cases reported in 2012 for a risk-adjusted rate of 11.1 per 1,000 elective surgery discharges. Eight hospitals had risk-adjusted rates that were statistically significantly higher than the statewide rate (see Table 6). Note that higher rate implies worse performance.
- Statewide, there were 62 '*Postoperative wound dehiscence*' cases reported in 2012 for a risk-adjusted rate of 1.7 per 1,000 abdominopelvic surgical discharges.
- There were 969 cases of '*Accidental punctures or lacerations*' reported by all 72 hospitals in 2012. The statewide risk-adjusted rate for occurrence of accidental puncture or laceration during procedure for the year was 1.2 per 1,000 discharges. Hospital-specific rates for this indicator ranged from a low of 0.0 to a high of 3.3 per 1,000 discharges.
- There were 170 cases of '*Birth trauma - injury to neonate*' reported

statewide in 2012 for a rate of 1.7 per 1,000 live births. Similarly, there were 473 cases of obstetric trauma among instrument-assisted vaginal deliveries (for a rate of 126.0 per 1,000 deliveries) and 1,091 cases of obstetric trauma among vaginal deliveries without instrument (for a rate of 19.1 per 1,000).

- Compared to the 2010 national estimates for PSIs (see Table 12), New Jersey appeared to have rates that were better than the national average for 7 of the 10 PSIs that are measured using rates. New Jersey rates were higher than the national rates only for 'Perioperative hemorrhage or hematoma' and 'Perioperative pulmonary embolism or deep vein thrombosis'. These differences may in part be, due to differences in years of data used and/or differences in data reporting by states.

Introduction

The Office of Health Care Quality Assessment (HCQA) of the New Jersey Department of Health and Senior Services (NJDHSS) assesses health care quality using qualitative and quantitative data reported by hospitals to support performance monitoring related to patient care and safety. Specifically, HCQA produces consumer reports on cardiac surgery, hospital performance, and hospital quality indicators; reviews confidential reports and root-cause analyses of reportable medical errors; and maintains several databases to support licensure requirements. In an effort to enhance the information the Department provides to the public on hospital care, HCQA staff apply statistical tools developed by the Federal Agency for Healthcare Research and Quality (AHRQ) to the New Jersey hospital discharge data commonly known as UB (Uniform Billing) data.

The AHRQ Quality Indicators (QIs) are a set of quality indicators organized into four modules, each of which measures quality associated, by and large, with patient care in an outpatient or inpatient setting. These four modules are: Prevention Quality Indicators (PQIs); Inpatient Quality Indicators (IQIs); Patient Safety Indicators (PSIs); and Pediatric Quality Indicators (PDIs). Background information on the development of these modules and the primary purposes they are designed to serve can be found at: www.nj.gov/health/healthcarequality/qi.shtml.

This report, presents findings resulting from the application of the Patient Safety Indicator (PSI) module to the 2012 New Jersey hospital discharge (UB) data. The PSI module contains 18 hospital-level indicators that reflect the quality of care provided by hospitals. These indicators serve as flags for potential quality problems (adverse events) rather than provide definitive measures of quality of care. According to the AHRQ, these indicators are selected based on their ability to screen out conditions present on admission from conditions that develop after admission, the potential preventability of the complication, and the ability of the indicator to identify medical error.

This report is a supplement to the Hospital Performance Report and covers only the 12 PSIs mandated for public reporting by law. In 2009, legislation (S2471) was signed into law requiring that hospital-specific data on patient-safety performance, and serious medical errors be included in the annual New Jersey Hospital Performance Report. Description of the Patient Safety Indicators module, Interpretation of the PSI measures as well as definitions of individual indicators presented in subsequent sections are, for the most part, excerpted from AHRQ's Guide and Software Documentation to Patient Safety Indicators. These sources are listed in the reference section of this report.

The tables present adverse events during hospitalization in each of the hospitals in the state. Risk-adjusted rates are provided along with confidence intervals for seven PSIs to help make a statistical assessment of patient safety in the hospital. Only observed rates are reported for the three birth delivery related PSIs because the

module does not risk-adjust these indicators. Two indicators – ‘foreign body left during procedure’ and ‘transfusion reaction’ - are reported in volume only because they are very rare events, commonly referred to as ‘never-events’. Statewide and national estimates are also provided to help compare hospital performance to the statewide or to the national average rates.

Comparison of a hospital’s rate to the statewide rate (presented in the top row of each of the PSI tables [Table 1-Table 11]) is one way to assess how well that hospital performed among its peers in the state. A hospital’s peers could be defined at many levels (e.g., teaching hospitals, urban hospitals, suburban hospitals, etc.). It is suggested that a hospital’s performance be assessed by looking at its performance across the several PSI estimates presented in the PSI tables.

The Patient Safety Indicators (PSIs) Module

Patient safety has been an issue of major national interest. Policy makers, providers, and consumers have made the safety of health care in U.S. hospitals a top priority. AHRQ states that the need to assess, monitor, track, and improve the safety of inpatient care became apparent with the publication of the Institute of Medicine’s series of reports describing the problems of medical errors.

One way of detecting and reporting potentially preventable adverse events is to develop screening measures based on routinely collected UB data. UB data provide adequate information (data elements) about health care services delivered in hospitals on patients’ diagnoses, procedures, age, gender, admission source, and discharge status. From these data elements, it is possible to construct a picture of the quality and safety of health care. Although quality assessments based on UB data cannot be definitive, they can be used to flag potential safety problems and success stories, which can then be further investigated. UB data can be used to identify indicators of potential problems that result from exposure to the health care system and are likely to be prevented as a result of system-level changes.

With this background, AHRQ developed the Patient Safety Indicators (PSIs) module in an effort to assess the quality of care inside hospitals with a focus on potentially preventable and other iatrogenic events, resulting from exposure to the health care system. The Patient Safety Indicators (PSIs) module is a tool specifically designed to help health care system leaders identify potential adverse events occurring during hospitalization for surgeries, procedures and childbirth. The PSIs (listed below) were developed after a comprehensive literature review, analysis of the International Classification of Diseases, 9th Revision, Clinical Modification, (ICD-9-CM) codes, review by a clinician panel, implementation of risk adjustment, and empirical analyses.

- Death in low mortality DRGs (PSI.02)
- Pressure ulcer (PSI.03)

- Death among surgical inpatients (PSI.04)
- Retained surgical item or unretrieved device fragment (PSI.05)*
- Foreign body left in during procedure (PSI.05)*
- Iatrogenic pneumothorax (PSI.06)*
- Central venous catheter-related bloodstream infections (PSI.07)
- Postoperative hip fracture (PSI.08)*
- Perioperative hemorrhage or hematoma Rate (PSI.09)*
- Postoperative physiologic and metabolic derangements (PSI.10)
- Postoperative respiratory failure (PSI.11)
- Perioperative pulmonary embolism or deep vein thrombosis (PSI.12)*
- Postoperative sepsis (PSI.13)*
- Postoperative wound dehiscence (PSI.14)*
- Accidental puncture and laceration (PSI.15)*
- Transfusion reaction (PSI.16)*
- Birth trauma - injury to neonate (PSI.17)*
- Obstetric trauma - vaginal delivery with instrument (PSI.18)*
- Obstetric trauma - vaginal delivery without instrument (PSI.19)*

The indicators have been shown to have complication/adverse event rates that vary substantially across institutions and for which evidence suggests that high complication/adverse event rates may be associated with deficiencies in the quality of care.

It is important to note that PSIs are intended to measure the occurrence rate of adverse events from: i) complications of medical conditions after admission, ii) complications from surgical procedures, and iii) complications from obstetric procedures. Fifteen of the 18 PSIs are related to surgical or medical discharges while the remaining three are for obstetric discharges. Six indicators (PSIs 03, 09, 10, 11, 12, 14), incorporate information about when procedures were performed (relative to the admission date), which is important in the risk-adjustment process. Admission type is used by four PSIs (PSIs 10, 11, 13, and 17) to identify elective surgeries and newborn admissions.

As stated earlier, this report focuses on the 12 PSIs (denoted by an asterisk in the list above) mandated for public reporting by the New Jersey legislature and provides comprehensive definitions for each along with their specific qualifications for their inclusion in the module.

How are PSI rates calculated?

The PSIs software module generates observed, expected, and risk-adjusted rates, as well as lower and upper 95% confidence limits for risk-adjusted rates, when applicable, for each indicator at a hospital level. Observed rates are the raw rates, while the expected and risk-adjusted rates are rates derived from applying the

average case-mix of a baseline file that reflects a large proportion of the U.S. hospitalized population.

Observed rates: Observed rates are raw rates generated by the software from the data under analysis. An observed rate is defined as the number of events of interest (numerator) divided by the population at risk (denominator). For hospital-level observed rates, the populations at risk are derived from hospital discharge records. The AHRQ software program calculates observed PSI rates regardless of the number of cases available. It is recommended that performance measurement assessment based on fewer than 30 cases in the denominator should be interpreted with caution.

Expected rates: Unlike observed rates, expected rates are derived from applying the average case-mix of a reference population that reflects a large proportion of the U.S. hospitalized population. The expected rate is the rate a hospital would have if it performed the same as the reference population, given the hospital's actual case-mix (e.g., age, gender, modified DRG and comorbidities).

Risk-adjusted rates: A hospital's risk-adjusted rate is obtained after its observed rate is adjusted to account for the difference between the patient case-mix of the reference population and that of the hospital. Regression coefficients from a baseline database reflecting a large proportion of the U.S. population (based on State Inpatient Databases (SID) compiled from about 44 states) are applied to observed rates for the purpose of making risk-adjustments. The baseline file of regression coefficients representing the average case-mix of the U.S. population is provided as part of the PSI software. The risk-adjusted rates reflect the age, sex, DRG, and comorbidity distribution of the data in the baseline file rather than the distributions of patients in the users' data. Thus, the observed rate (raw indicator) is adjusted using a logistic regression to account for differences among hospitals and areas in demographics. This will allow risk-adjusted rates produced by various states to be compared directly to one another. The interpretation of risk-adjusted rates becomes clear when we compare risk-adjusted rates with the observed rates. Hospitals that exhibit large differences between their observed and risk-adjusted rates tend to have a more complex case-mix.

Risk-adjustment in the latter version of the PSI module includes an adjustment for the Present on Admission (POA) indicator. The POA indicator identifies instances in which a condition was present on admission (i.e. pre-existing condition) and those that occur during the hospital stay. The POA indicator enables conditions present on admission to be identified and excluded from the quality measures, when appropriate.

Interpretation of PSI rates

The purpose of the analysis determines which rates one should use in evaluating the performance of a hospital. If the user's primary interest is to focus on a particular hospital, to identify cases for further follow-up and quality improvement without

comparisons made to other hospitals, then he/she ought to simply examine the observed rate. But, if the purpose of the analysis is to compare the performance of a particular hospital with national, state, or regional averages or performances of other selected hospitals, then all rates (observed, expected and risk-adjusted) should be examined.

Hospitals can compare their expected rates to the population rate to see how their patient case-mix compares to the reference population. The population rate refers to the overall rate for the reference population. If the population rate is higher than the expected rate, then the hospital's case-mix is less severe than the reference population. If the population rate is lower than the expected rate, then the hospital's case-mix is more severe than the reference population.

Comparing the observed rate to the expected rate allows hospitals to see how far or how close they are from what is expected of them, based on the reference population. If the observed rate is higher than the expected rate for any given indicator (i.e., the ratio of observed/expected is greater than 1.0), then the implication is that the hospital performed worse than expected for that particular indicator. If the observed rate is lower than the expected rate (i.e., the ratio of observed/expected is less than 1.0), then the implication is that the hospital performed better than expected.

Comparing a hospital's risk-adjusted rate to its expected rate shows the effect of risk-adjustment on the patient safety indicator measurement. The risk-adjusted rate accounts for the difference between the case-mix of the reference population and the hospital's case-mix. For that reason, risk-adjusted rates should be used for better hospital-to-hospital comparisons. Risk-adjusted rates are given along with their respective 95% confidence intervals.

- Even in the best hospital, some patients will experience complications either after an operation or as a result of other care. The rates in this report are calculated by comparing the number of complications (adverse events) expected in a particular hospital (based on the number of operations they do or patients they see, usually after adjusting for how old and how sick their patients are) and how many patients actually experienced the adverse events (complications).
- Confidence Intervals are used to identify which hospitals had statistically significantly more or fewer complications than expected given the risk factors of their patients. Hospitals with significantly higher rates than expected, after adjusting for risk factors, are those where the confidence interval range falls entirely above the statewide risk-adjusted complication rate. Hospitals with statistically significantly lower rates than expected have their confidence interval range entirely below the statewide risk-adjusted complication rate.
- Comparison of a hospital's rate to the statewide rate (presented in the top row of each PSI Table), is one way to assess how well that hospital performed

among its peers in the state. A hospital's peers could be defined at many levels (e.g., teaching hospitals, urban hospitals, suburban hospitals, etc.). It is suggested that a hospital's performance be assessed by looking at its performance across the 12 PSI estimates presented in the Tables.

A hospital's rate is statistically significantly above (denoted by double asterisks) the statewide rate if the statewide rate falls completely below the hospital's 95% confidence interval. By comparison, a hospital's rate is significantly below (denoted by a single asterisk) the statewide rate if the statewide rate falls completely above the hospital confidence interval for that indicator. Some rates that appear large are not marked as statistically significantly higher than the statewide rate while others that appear small are not marked as statistically lower than the statewide rate. The reason may be that rates calculated from small numbers of events tend to have wider confidence intervals that make the statewide rate fall within the interval, giving the appearance of good performance by those hospitals compared to hospitals with rates based on large numbers of events. For example, the risk-adjusted rate for Perioperative Hemorrhage or Hematoma (Table 4) for Hackensack University Medical Center is 18.6 per 1,000 surgical discharges. This rate of 18.6 per 1,000 which is derived from 70 cases out of 10,003 surgical discharges has a 95% confidence interval of 16.2 – 21.0 and is considered statistically significantly higher than the statewide rate of 14.7 per 1,000 because the confidence interval range is completely above the statewide average. By comparison, Mountainside Hospital's rate of 19.1 per 1,000, which is derived from 14 cases out of 1,787 surgeries has a 95% confidence interval of 13.6 – 24.5 and is not statistically significantly different from the statewide rate of 14.7 per 1,000 because the statewide average falls within the hospital's confidence interval range.

In general, PSIs are not intended as definitive quality measures because quality of performance may be influenced by several other factors. However, there is strong evidence that PSI measures indicate differences in hospital performance, which are potentially clinically important. They do measure differences in the hospitals' ability to reduce severe and potentially preventable complications and adverse events. Performance on a single PSI often cannot reliably indicate actual quality differences. AHRQ recommends that examining all the indicators together is likely to produce a more complete picture of overall quality of care.

Limitations of PSI measures

These PSI rates should only be seen as a starting point for examining the quality of care at a particular hospital. They should not be used to make strong conclusions. There are a few issues to keep in mind when looking at these measures.

- The PSIs do not address all aspects of quality. For example, they do not include information on what patients say about their care in the hospital, or information on whether hospitals consistently follow steps known to lead to better results.

- In some cases, the PSIs track serious failures in a hospital's performance which happen only once in a while. One has to be careful when comparing hospitals on these very rare events. The numbers are so small that it is difficult to know when a difference is meaningful or occurs due to chance alone. For example, a major reaction to a blood transfusion occurs in only a few cases out of a million people each year.
- One obvious limitation is that many important quality concerns including adverse drug events cannot be monitored using UB data because these data are unlikely to capture all cases of patient complications. The indicators in the PSIs module contain a large proportion of surgical indicators rather than medical or psychiatric indicators because medical complications are often difficult to distinguish from comorbidities that are present on admission. In addition, patients with medical conditions tend to be more heterogeneous than surgical patients, especially elective surgical patients, making it more difficult to account for case-mix.
- Incomplete reporting is an issue in the accuracy of any data source used for identifying patient safety problems, as medical providers might fear adverse consequences as a result of "full disclosure" in potentially public records such as discharge abstracts.
- The ability of administrative data to distinguish between adverse events in which no error occurred from true medical errors is limited. A number of factors such as heterogeneity of clinical conditions included in some codes, lack of information about event timing available in these data sets, and limited clinical detail for risk adjustment, contribute to the difficulty in identifying complications that represent medical error or may at least be in some part preventable.
- Questions about the clinical accuracy of discharge-based diagnosis coding lead to concerns about the interpretation of reported diagnoses that may represent safety problems. Specifically, UB data are unlikely to capture all cases of a complication, regardless of the preventability, without false positives and false negatives (sensitivity and specificity). Also, when the codes are accurate in defining an event, the clinical vagueness inherent in the description of the code itself (e.g., "hypotension") may lead to a highly heterogeneous pool of clinical states represented by that code.

Patient Safety Indicator estimates for New Jersey

As indicated earlier, this report is based on an application of the AHRQ PSIs module to the New Jersey hospital discharge data. In this section, we provide an abbreviated description or definition for each of the 12 indicators used, followed by a table showing the numbers of adverse events, total hospital discharges, and the

corresponding observed, expected and risk-adjusted rates, along with 95% confidence intervals, when applicable. Where the cell entry is missing, it is designated by “.” to indicate that the hospital did not perform that particular procedure or it did less than 3 procedures (risk-adjusted rate is not computed when the denominator is less than 3).

Comparison of a specific hospital-level PSI rate to the statewide average for the same indicator is one appropriate way to see how well a hospital performs among its peers. Following AHRQ’s recommendation, we have compared hospital rates against statewide rates to assess performance.

PSI.05 - Retained surgical item or unretrieved device fragment

- This indicator is measured using volume of occurrence – not a rate. It tells you the number of patients who had a ‘retained surgical item or unretrieved device fragment’ during surgical or medical procedures. It is considered a never-event and happens very rarely. All cases with pre-existing conditions are excluded from the measure.
- The measure refers to discharges 18 years and older or obstetric patients (MDC 14 - pregnancy, childbirth, and puerperium), with ICD-9-CM codes for a ‘retained surgical item or unretrieved device fragment’ in any secondary diagnosis field of medical and surgical discharges defined by specific DRGs or MS-DRGs.
- Patients with ICD-9-CM codes for a ‘retained surgical item or unretrieved device fragment’ in the principal diagnosis field or secondary diagnosis present on admission are excluded from the measure.
- Table 1 shows that, Statewide there were 38 cases of ‘retained surgical item or unretrieved device fragment’ in 2012.

Table 1. Retained Surgical Item or Unretrieved Device Fragment (formerly Foreign body left during procedure)

Hospital	# of cases
Statewide	38
Atlanticare Regional Medical Center-City	1
Bergen Regional Medical Center	1
Christ Hospital	1
Clara Maass Medical Center	1
Cooper Hospital/University Medical Center	1
Hackensack University Medical Center	11
Hoboken University Medical Center	1
Holy Name Medical Center	1
Jersey Shore University Medical Center	2
Kennedy University Hospital - Wash. Twp.	1
Morristown Memmorial Hospital	1
Mountainside Hospital	1
Newark Beth Israel Medical Center	1
Raritan Bay Medical Center-Old Bridge	1
RWJ University Hospital	3
RWJ University Hospital at Hamilton	1
St. Mary's Hospital (Passaic)	1
UMDNJ-University Hospital	3
Valley Hospital	2
Virtua-Memorial Hospital Burlington Cty.	2
Virtua-West Jersey Hospital Voorhees	1

Source: New Jersey 2012 UB Data.

PSI.06 - Iatrogenic pneumothorax

- This indicator flags cases of iatrogenic pneumothorax (i.e. patients who had air leaking out of their lungs due to an accidental puncture during a medical or surgical procedure) occurring in a facility. The rate, which is risk-adjusted by age, sex, DRG, and comorbidity categories, is defined as the number of iatrogenic pneumothorax cases per 1,000 discharges surgical and medical discharges.
- The numerator refers to the number of discharges with ICD-9-CM code of 512.1 in any secondary diagnosis field among cases meeting the inclusion and exclusion rules for the denominator.
- The denominator refers to all medical and surgical discharges age 18 years and older defined by specific DRGs.
- The following cases are excluded from the denominator or from the rate calculation:
 - cases with ICD-9-CM code of 512.1 as the principal diagnosis or secondary diagnosis present on admission, if known;
 - cases with ICD-9-CM diagnosis code of chest trauma or pleural effusion;
 - cases with ICD-9-CM procedure code of diaphragmatic surgery repair;
 - cases with any code indicating thoracic surgery or lung or pleural biopsy or assigned to cardiac surgery DRGs; and
 - obstetric cases (MDC 14 - pregnancy, childbirth, and puerperium).
- Table 2 shows the number of iatrogenic pneumothorax cases by hospital, as well as the observed, expected and risk-adjusted rates with their corresponding 95% confidence intervals. Statewide, there were 238 cases of iatrogenic pneumothorax out of 700,999 eligible discharges reported in 2012 for a risk-adjusted rate of 0.3 per 1,000 discharges.

Table 2. Iatrogenic Pneumothorax (per 1,000 medical and surgical discharges)

Hospital	# of cases	# of discharges	Obs. rate	Exp. Rate Δ	Risk-adjusted rate	95% Confidence interval
Statewide	238	700,999	0.3	0.5	0.3	0.3 - 0.4
Atlanticare Regional Medical Center-City	1	8,724	0.1	0.4	0.1	0.0 - 0.6
Atlanticare Regional Medical Center-Mainland	1	14,862	0.1	0.4	0.1	0.0 - 0.4
Bayonne Medical Center	1	4,963	0.2	0.4	0.2	0.0 - 0.8
Bayshore Community Hospital	1	6,727	0.1	0.5	0.1	0.0 - 0.6
Bergen Regional Medical Center	0	6,096	0.0	0.4	0.0	0.0 - 0.6
Cape Regional Medical Center	1	7,999	0.1	0.3	0.2	0.0 - 0.7
Capital Health Medical Center - Hopewell	3	7,246	0.4	0.4	0.5	0.0 - 1.0
Capital Health Regional Medical Center	2	9,250	0.2	0.4	0.3	0.0 - 0.7
CentraState Medical Center	1	10,766	0.1	0.4	0.1	0.0 - 0.5
Chilton Memorial Hospital	6	7,866	0.8	0.4	0.9 **	0.4 - 1.4
Christ Hospital	1	6,922	0.1	0.4	0.2	0.0 - 0.7
Clara Maass Medical Center	6	13,629	0.4	0.4	0.4	0.1 - 0.8
Community Medical Center	2	21,946	0.1	0.5	0.1	0.0 - 0.4
Cooper Hospital/University Medical Center	12	17,223	0.7	0.5	0.6	0.3 - 0.9
Deborah Heart and Lung Center	2	3,908	0.5	1.4	0.2	0.0 - 0.5
East Orange General Hospital	1	7,020	0.1	0.5	0.1	0.0 - 0.6
Englewood Hospital and Medical Center	1	12,294	0.1	0.5	0.1	0.0 - 0.4
Hackensack University Medical Center	5	25,854	0.2	0.5	0.2	0.0 - 0.4
Hackettstown Regional Medical Center	1	3,635	0.3	0.4	0.3	0.0 - 1.0
Hoboken University Medical Center	2	4,295	0.5	0.3	0.6	0.0 - 1.3
Holy Name Medical Center	4	10,855	0.4	0.5	0.3	0.0 - 0.7
Hunterdon Medical Center	8	5,973	1.3	0.4	1.4 **	0.8 - 1.9
Jersey City Medical Center	3	10,843	0.3	0.4	0.3	0.0 - 0.7
Jersey Shore University Medical Center	10	17,198	0.6	0.7	0.4	0.1 - 0.6
JFK Medical Center/Anthony M. Yelencis	2	13,306	0.2	0.5	0.1	0.0 - 0.4
Kennedy University Hospital - Cherry Hill	1	6,899	0.1	0.3	0.2	0.0 - 0.8
Kennedy University Hospital - Stratford	0	6,561	0.0	0.3	0.0	0.0 - 0.6
Kennedy University Hospital - Wash. Twp.	5	12,119	0.4	0.4	0.5	0.1 - 0.9
Kimball Medical Center	1	8,372	0.1	0.3	0.2	0.0 - 0.7
Lourdes Medical Center of Burlington Cty.	2	6,462	0.3	0.3	0.4	0.0 - 1.0
Meadowlands Hospital Medical Center	1	2,076	0.5	0.5	0.5	0.0 - 1.3
Memorial Hospital of Salem County	1	3,485	0.3	0.3	0.4	0.0 - 1.2
Monmouth Medical Center	0	9,026	0.0	0.4	0.0	0.0 - 0.5
Morristown Memorial Hospital	2	22,114	0.1	0.5	0.1	0.0 - 0.3
Mountainside Hospital	2	7,596	0.3	0.5	0.3	0.0 - 0.7
Newark Beth Israel Medical Center	13	12,743	1.0	0.5	0.8 **	0.5 - 1.2
Newton Medical Center	2	6,050	0.3	0.4	0.4	0.0 - 1.0
Ocean Medical Center - Bricktown	3	11,512	0.3	0.4	0.3	0.0 - 0.6
Our Lady of Lourdes Medical Center	4	10,200	0.4	0.6	0.3	0.0 - 0.6
Overlook Medical Center	4	15,917	0.3	0.5	0.2	0.0 - 0.6
Palisades Medical Center - NY PHS	1	7,424	0.1	0.4	0.2	0.0 - 0.7
Raritan Bay Medical Center-Old Bridge	0	4,413	0.0	0.5	0.0	0.0 - 0.6
Raritan Bay Medical Center-Perth Amboy	4	6,654	0.6	0.4	0.6	0.1 - 1.1

Table 2. Iatrogenic Pneumothorax (per 1,000 medical and surgical discharges)

Hospital	# of cases	# of discharges	Obs. rate	Exp. Rate Δ	Risk-adjusted rate	95% Confidence interval
Statewide	238	700,999	0.3	0.5	0.3	0.3 - 0.4
Riverview Medical Center	4	9,222	0.4	0.5	0.4	0.0 - 0.8
RWJ University Hospital	19	23,557	0.8	0.6	0.6 **	0.4 - 0.8
RWJ University Hospital at Hamilton	4	9,516	0.4	0.4	0.5	0.0 - 1.0
RWJ University Hospital at Rahway	3	6,138	0.5	0.7	0.3	0.0 - 0.7
Shore Medical Center	3	8,529	0.4	0.4	0.4	0.0 - 0.9
Somerset Medical Center	2	11,774	0.2	0.5	0.2	0.0 - 0.5
South Jersey Healthcare Regional MC	8	12,396	0.6	0.4	0.7	0.3 - 1.1
South Jersey Hospital-Elmer	1	3,249	0.3	0.3	0.4	0.0 - 1.2
Southern Ocean Medical Center	0	5,131	0.0	0.4	0.0	0.0 - 0.6
St. Barnabas Medical Center	8	17,155	0.5	0.5	0.4	0.1 - 0.7
St. Clare's Hospital-Denville	1	9,459	0.1	0.4	0.1	0.0 - 0.6
St. Clare's Hospital-Dover	0	3,566	0.0	0.4	0.0	0.0 - 0.7
St. Clare's Hospital-Sussex	0	550	0.0	0.2	0.0	0.0 - 2.5
St. Francis Medical Center-Trenton	3	5,926	0.5	0.4	0.6	0.0 - 1.1
St. Joseph's Hospital and Medical Center	11	20,502	0.5	0.4	0.6	0.3 - 0.9
St. Joseph's Wayne Hospital	5	5,205	1.0	0.5	0.9	0.3 - 1.4
St. Luke Warren Hospital	0	4,307	0.0	0.4	0.0	0.0 - 0.7
St. Mary's Hospital (Passaic)	6	6,907	0.9	0.6	0.7	0.2 - 1.1
St. Michael's Medical Center	4	8,666	0.5	0.5	0.4	0.0 - 0.8
St. Peter's University Hospital	0	9,398	0.0	0.5	0.0	0.0 - 0.4
Trinitas Regional Medical Center	3	9,970	0.3	0.4	0.3	0.0 - 0.8
UMDNJ-University Hospital	11	11,726	0.9	0.5	0.8 **	0.5 - 1.2
Underwood-Memorial Hospital	7	9,674	0.7	0.4	0.8 **	0.4 - 1.3
University Medical Center at Princeton	4	9,623	0.4	0.5	0.4	0.0 - 0.8
Valley Hospital	4	20,317	0.2	0.5	0.2	0.0 - 0.5
Virtua-Memorial Hospital Burlington Cty.	3	13,454	0.2	0.5	0.2	0.0 - 0.6
Virtua-West Jersey Hospital Berlin	0	3,471	0.0	0.5	0.0	0.0 - 0.7
Virtua-West Jersey Hospital Marlton	1	9,966	0.1	0.5	0.1	0.0 - 0.5
Virtua-West Jersey Hospital Voorhees	0	12,622	0.0	0.5	0.0	0.0 - 0.3

Source: New Jersey 2012 UB Data.

** Statistically significantly above statewide average (i.e., worse than statewide average).

Δ : Expected rate is the rate the hospital would have if it had the same case-mix (e.g., age, gender, DRG, and comorbidity categories) as the reference or statewide population. If the observed rate is higher than the expected rate (i.e., the ratio of observed to expected is greater than 1.0), it suggests that the hospital performed worse than the reference population on that indicator.

PSI.08 – Postoperative hip fracture

- This indicator intends to capture cases of in-hospital hip fractures and includes only secondary diagnosis codes to eliminate fractures that were present on admission. The rate, which is risk-adjusted for age, sex, DRG, and comorbidity categories, is defined as the number of cases of in-hospital hip fracture per 1,000 surgical discharges with an operating room procedure.
- The numerator refers to discharges with ICD-9-CM code for hip fracture in any secondary diagnosis field among cases meeting the inclusion and exclusion rules for the denominator.
- The denominator refers to all surgical discharges 18 years and older defined by specific DRGs and an ICD-9-CM code for an operating room procedure.
- The following cases are excluded from the denominator or from rate calculation:
 - cases with ICD-9-CM code for hip fracture in the principal diagnosis field or secondary diagnosis present on admission, if known;
 - cases where the only operating room procedure is hip fracture repair; and where a procedure for hip fracture repair occurs before or on the same day as the first operating room procedure (*if day of procedure is not available in the input data file, the rate may be slightly lower than if the information was available*);
 - cases with diseases and disorders of the musculoskeletal system and connective tissue (MDC 8);
 - cases with principal diagnosis or secondary diagnosis (present on admission, if known) of seizure, syncope, stroke, coma, cardiac arrest, poisoning, trauma, delirium and other psychoses, or anoxic brain injury;
 - cases with any diagnosis of metastatic cancer, lymphoid malignancy or bone malignancy, or self-inflicted injury; and
 - MDC 14 (pregnancy, childbirth and the puerperium).
- Table 3 shows the number, by hospital, of discharges with postoperative hip fracture among all surgical discharges age 18 and older, the observed rates, expected rates and risk-adjusted rates with their corresponding 95% confidence intervals. Statewide, there were only 3 postoperative hip fracture cases reported out of 126,198 eligible discharges in 2012.

Table 3. Postoperative Hip Fracture (per 1,000 surgical discharges)

Hospital	# of cases	# of discharges	Obs. rate	Exp. Rate Δ	Risk-adjusted rate	95% Confidence interval
Statewide	3	126,198	0.0	0.0	0.0	0.0 - 0.1
Atlanticare Regional Medical Center-City	0	1,094	0.0	0.0	0.0	0.0 - 0.4
Atlanticare Regional Medical Center-Mainland	0	2,374	0.0	0.0	0.0	0.0 - 0.2
Bayonne Medical Center	0	600	0.0	0.0	0.0	0.0 - 0.5
Bayshore Community Hospital	0	720	0.0	0.0	0.0	0.0 - 0.4
Bergen Regional Medical Center	0	110	0.0	0.0	0.0	0.0 - 1.1
Cape Regional Medical Center	0	673	0.0	0.0	0.0	0.0 - 0.5
Capital Health Medical Center - Hopewell	0	1,471	0.0	0.0	0.0	0.0 - 0.3
Capital Health Regional Medical Center	1	691	1.4	0.0	1.6	0.0 - 2.1
CentraState Medical Center	0	1,767	0.0	0.0	0.0	0.0 - 0.3
Chilton Memorial Hospital	0	1,174	0.0	0.0	0.0	0.0 - 0.3
Christ Hospital	0	984	0.0	0.0	0.0	0.0 - 0.4
Clara Maass Medical Center	0	2,852	0.0	0.0	0.0	0.0 - 0.2
Community Medical Center	0	2,950	0.0	0.0	0.0	0.0 - 0.2
Cooper Hospital/University Medical Center	0	4,613	0.0	0.0	0.0	0.0 - 0.2
Deborah Heart and Lung Center	0	2,767	0.0	0.0	0.0	0.0 - 0.2
East Orange General Hospital	0	430	0.0	0.0	0.0	0.0 - 0.6
Englewood Hospital and Medical Center	0	3,519	0.0	0.0	0.0	0.0 - 0.2
Hackensack University Medical Center	0	6,203	0.0	0.0	0.0	0.0 - 0.1
Hackettstown Regional Medical Center	0	437	0.0	0.0	0.0	0.0 - 0.6
Hoboken University Medical Center	0	588	0.0	0.0	0.0	0.0 - 0.5
Holy Name Medical Center	0	1,965	0.0	0.0	0.0	0.0 - 0.3
Hunterdon Medical Center	0	747	0.0	0.0	0.0	0.0 - 0.4
Jersey City Medical Center	0	1,485	0.0	0.0	0.0	0.0 - 0.3
Jersey Shore University Medical Center	0	4,287	0.0	0.0	0.0	0.0 - 0.2
JFK Medical Center/Anthony M. Yelensics	0	2,413	0.0	0.0	0.0	0.0 - 0.2
Kennedy University Hospital - Cherry Hill	0	326	0.0	0.0	0.0	0.0 - 0.6
Kennedy University Hospital - Stratford	0	936	0.0	0.0	0.0	0.0 - 0.4
Kennedy University Hospital - Wash. Twp.	0	1,436	0.0	0.0	0.0	0.0 - 0.3
Kimball Medical Center	0	566	0.0	0.0	0.0	0.0 - 0.5
Lourdes Medical Center of Burlington Cty.	0	852	0.0	0.0	0.0	0.0 - 0.4
Meadowlands Hospital Medical Center	0	346	0.0	0.0	0.0	0.0 - 0.6
Memorial Hospital of Salem County	0	394	0.0	0.0	0.0	0.0 - 0.6
Monmouth Medical Center	0	1,501	0.0	0.0	0.0	0.0 - 0.3
Morristown Memorial Hospital	0	6,941	0.0	0.0	0.0	0.0 - 0.1
Mountainside Hospital	0	1,307	0.0	0.0	0.0	0.0 - 0.3
Newark Beth Israel Medical Center	1	3,128	0.3	0.0	0.3	0.0 - 0.5
Newton Medical Center	0	582	0.0	0.0	0.0	0.0 - 0.5
Ocean Medical Center - Bricktown	0	1,564	0.0	0.0	0.0	0.0 - 0.3
Our Lady of Lourdes Medical Center	0	3,567	0.0	0.0	0.0	0.0 - 0.2
Overlook Medical Center	0	3,324	0.0	0.0	0.0	0.0 - 0.2
Palisades Medical Center - NY PHS	0	878	0.0	0.0	0.0	0.0 - 0.4
Raritan Bay Medical Center-Old Bridge	0	521	0.0	0.0	0.0	0.0 - 0.5

Table 3. Postoperative Hip Fracture (per 1,000 surgical discharges)

Hospital	# of cases	# of discharges	Obs. rate	Exp. Rate Δ	Risk-adjusted rate	95% Confidence interval
Statewide	3	126,198	0.0	0.0	0.0	0.0 - 0.1
Raritan Bay Medical Center-Perth Amboy	0	883	0.0	0.0	0.0	0.0 - 0.4
Riverview Medical Center	0	1,660	0.0	0.0	0.0	0.0 - 0.3
RWJ University Hospital	0	6,301	0.0	0.0	0.0	0.0 - 0.1
RWJ University Hospital at Hamilton	0	1,443	0.0	0.0	0.0	0.0 - 0.3
RWJ University Hospital at Rahway	0	639	0.0	0.0	0.0	0.0 - 0.5
Shore Medical Center	0	963	0.0	0.0	0.0	0.0 - 0.4
Somerset Medical Center	0	1,892	0.0	0.0	0.0	0.0 - 0.3
South Jersey Healthcare Regional MC	0	1,917	0.0	0.0	0.0	0.0 - 0.3
South Jersey Hospital-Elmer	0	366	0.0	0.0	0.0	0.0 - 0.6
Southern Ocean Medical Center	0	644	0.0	0.0	0.0	0.0 - 0.5
St. Barnabas Medical Center	0	4,549	0.0	0.0	0.0	0.0 - 0.2
St. Clare's Hospital-Denville	0	1,150	0.0	0.0	0.0	0.0 - 0.3
St. Clare's Hospital-Dover	0	465	0.0	0.0	0.0	0.0 - 0.5
St. Clare's Hospital-Sussex	0	38	0.0	0.0	0.0	0.0 - 1.9
St. Francis Medical Center-Trenton	0	1,104	0.0	0.0	0.0	0.0 - 0.3
St. Joseph's Hospital and Medical Center	0	3,388	0.0	0.0	0.0	0.0 - 0.2
St. Joseph's Wayne Hospital	0	599	0.0	0.0	0.0	0.0 0.5
St. Luke Warren Hospital	0	504	0.0	0.0	0.0	0.0 - 0.5
St. Mary's Hospital (Passaic)	0	1,499	0.0	0.0	0.0	0.0 - 0.3
St. Michael's Medical Center	0	1,567	0.0	0.0	0.0	0.0 - 0.3
St. Peter's University Hospital	0	2,067	0.0	0.0	0.0	0.0 - 0.3
Trinitas Regional Medical Center	0	1,477	0.0	0.0	0.0	0.0 - 0.3
UMDNJ-University Hospital	0	2,006	0.0	0.0	0.0	0.0 - 0.3
Underwood-Memorial Hospital	0	1,316	0.0	0.0	0.0	0.0 - 0.3
University Medical Center at Princeton	0	1,484	0.0	0.0	0.0	0.0 - 0.3
Valley Hospital	0	4,383	0.0	0.0	0.0	0.0 - 0.2
Virtua-Memorial Hospital Burlington Cty.	0	2,606	0.0	0.0	0.0	0.0 - 0.2
Virtua-West Jersey Hospital Berlin	0	175	0.0	0.0	0.0	0.0 - 0.9
Virtua-West Jersey Hospital Marlton	0	1,724	0.0	0.0	0.0	0.0 - 0.3
Virtua-West Jersey Hospital Voorhees	1	2,306	0.4	0.0	0.5	0.0 - 0.7

Source: New Jersey 2012 UB Data.

Δ : Expected rate is the rate the hospital would have if it had the same case-mix (e.g., age, gender, DRG, and comorbidity categories) as the reference or statewide population. If the observed rate is higher than the expected rate (i.e., the ratio of observed to expected is greater than 1.0), it suggests that the hospital performed worse than the reference population on that indicator.

PSI.09 – Perioperative hemorrhage or hematoma

- This indicator is designed to capture perioperative hemorrhage or hematoma cases with control of perioperative hemorrhage, drainage of hematoma, or a miscellaneous hemorrhage- or hematoma-related procedure following surgery. The rate, which is risk-adjusted for age, sex, DRG, and comorbidity categories, is defined as the number of patients with perioperative hemorrhage (too much bleeding) or hematoma (drainage of hematoma) per 1,000 surgical discharges ages 18 years and older.
- The numerator refers to the number of discharges with ICD-9-CM codes for perioperative hemorrhage or hematoma in any secondary diagnosis field or discharges with ICD-9-CM codes for perioperative control of hemorrhage or drainage of hematoma in any secondary procedure, while the denominator refers to all surgical discharges age 18 years and older defined by specific DRGs and an ICD-9-CM code for an operating room procedure.
- The following cases are excluded from the denominator or from rate calculation:
 - cases with a secondary diagnosis of perioperative hemorrhage or hematoma present on admission, if known;
 - cases with a diagnosis of coagulation disorder;
 - cases with a principal diagnosis of perioperative hemorrhage or hematoma;
 - cases where the only operating room procedure is control of perioperative hemorrhage, drainage of hematoma, or a miscellaneous hemorrhage- or hematoma-related procedure;
 - cases where a procedure for perioperative control of hemorrhage or drainage of hematoma occurs before the first operating room procedure (*if day of procedure is not available in the input data file, the rate may be slightly lower than if the information was available*); and
 - obstetric cases (MDC 14 - pregnancy, childbirth and the puerperium).
- Table 4 shows the number of perioperative hemorrhage or hematoma cases by hospital, the number of eligible surgical discharges, observed rates, expected rates, and risk-adjusted rates along with their corresponding 95% confidence intervals. Statewide, there were 1,086 perioperative hemorrhage or hematoma cases out of 185,387 eligible surgical discharges reported in 2012 for a statewide risk-adjusted rate of 14.7 per 1,000.

Table 4. Perioperative Hemorrhage or Hematoma (per 1,000 surgical discharges)

Hospital	# of cases	# of discharges	Obs. rate	Exp. Rate Δ	Risk-adjusted rate	95% Confidence interval
Statewide	1,086	185,387	5.9	2.3	14.7	14.2 - 15.3
Atlanticare Regional Medical Center-City	5	1,765	2.8	2.2	7.3 *	1.6 - 13.0
Atlanticare Regional Medical Center-Mainland	11	4,745	2.3	1.5	9.1 *	4.8 - 13.3
Bayonne Medical Center	3	760	3.9	2.4	9.3	1.0 - 17.5
Bayshore Community Hospital	2	1,062	1.9	2.4	4.5 *	0.0 - 11.5
Bergen Regional Medical Center	1	167	6.0	2.1	16.3	0.0 - 35.2
Cape Regional Medical Center	5	1,171	4.3	2.1	11.7	4.5 - 18.9
Capital Health Medical Center - Hopewell	10	2,203	4.5	2.4	10.7	5.9 - 15.6
Capital Health Regional Medical Center	5	1,524	3.3	2.1	8.8	2.6 - 15.0
CentraState Medical Center	15	2,560	5.9	2.3	14.6	10.0 - 19.2
Chilton Memorial Hospital	19	1,775	10.7	2.4	25.9 **	20.4 - 31.4
Christ Hospital	7	1,127	6.2	2.7	13.2	6.8 - 19.7
Clara Maass Medical Center	19	3,515	5.4	2.4	12.9	9.0 - 16.7
Community Medical Center	23	4,400	5.2	2.3	13.0	9.5 - 16.5
Cooper Hospital/University Medical Center	35	6,631	5.3	2.7	11.4 *	8.7 - 14.1
Deborah Heart and Lung Center	18	2,757	6.5	2.6	14.4	10.2 - 18.6
East Orange General Hospital	3	525	5.7	2.7	12.2	2.8 - 21.7
Englewood Hospital and Medical Center	9	4,936	1.8	2.1	4.9 *	1.5 - 8.4
Hackensack University Medical Center	70	10,003	7.0	2.2	18.6 **	16.2 - 21.0
Hackettstown Regional Medical Center	2	770	2.6	2.0	7.5	0.0 - 16.5
Hoboken University Medical Center	0	832	0.0	2.1	0.0 *	0.0 - 8.5
Holy Name Medical Center	5	2,873	1.7	2.3	4.4 *	0.0 - 8.8
Hunterdon Medical Center	8	1,289	6.2	1.9	18.7	11.5 - 25.9
Jersey City Medical Center	12	2,005	6.0	2.2	15.5	10.2 - 20.9
Jersey Shore University Medical Center	59	6,362	9.3	2.2	24.2 **	21.2 - 27.2
JFK Medical Center/Anthony M. Yelencis	16	3,500	4.6	2.2	12.2	8.1 - 16.3
Kennedy University Hospital - Cherry Hill	0	517	0.0	2.0	0.0 *	0.0 - 10.9
Kennedy University Hospital - Stratford	4	1,105	3.6	2.2	9.3	2.1 - 16.4
Kennedy University Hospital - Wash. Twp.	4	2,537	1.6	2.0	4.6 *	0.0 - 9.6
Kimball Medical Center	2	758	2.6	2.5	6.1 *	0.0 - 14.2
Lourdes Medical Center of Burlington Cty.	5	1,141	4.4	2.3	11.0	4.0 - 17.9
Meadowlands Hospital Medical Center	0	466	0.0	2.3	0.0 *	0.0 - 10.8
Memorial Hospital of Salem County	1	484	2.1	2.2	5.4	0.0 - 16.3
Monmouth Medical Center	11	2,619	4.2	2.2	11.1	6.4 - 15.8
Morristown Memorial Hospital	116	11,163	10.4	2.1	28.5 **	26.2 - 30.8
Mountainside Hospital	14	1,787	7.8	2.4	19.1	13.6 - 24.5
Newark Beth Israel Medical Center	42	3,549	11.8	2.9	23.8 **	20.2 - 27.3
Newton Medical Center	4	961	4.2	2.1	11.1	3.3 - 19.0
Ocean Medical Center - Bricktown	15	2,949	5.1	2.0	14.3	9.7 - 18.9
Our Lady of Lourdes Medical Center	42	3,756	11.2	2.7	23.7 **	20.2 - 27.2
Overlook Medical Center	29	5,126	5.7	2.2	14.7	11.3 - 18.0
Palisades Medical Center - NY PHS	3	1,047	2.9	2.5	6.5 *	0.0 - 13.4
Raritan Bay Medical Center-Old Bridge	1	686	1.5	2.3	3.6 *	0.0 - 12.5
Raritan Bay Medical Center-Perth Amboy	3	1,088	2.8	2.1	7.4	0.0 - 14.7

Table 4. Perioperative Hemorrhage or Hematoma (per 1,000 surgical discharges)

Hospital	# of cases	# of discharges	Obs. rate	Exp. Rate Δ	Risk-adjusted rate	95% Confidence interval
Statewide	1,086	185,387	5.9	2.3	14.7	14.2 - 15.3
Riverview Medical Center	9	2,820	3.2	2.1	8.5 *	4.0 - 13.1
RWJ University Hospital	88	8,534	10.3	2.5	23.7 **	21.3 - 26.2
RWJ University Hospital at Hamilton	12	2,154	5.6	2.2	14.8	9.6 - 20.0
RWJ University Hospital at Rahway	0	902	0.0	2.5	0.0 *	0.0 - 7.5
Shore Medical Center	8	2,032	3.9	1.9	11.6	6.0 - 17.3
Somerset Medical Center	14	2,751	5.1	2.1	14.0	9.3 - 18.7
South Jersey Healthcare Regional MC	11	2,676	4.1	2.4	9.9 *	5.4 - 14.3
South Jersey Hospital-Elmer	1	726	1.4	1.7	4.6 *	0.0 - 14.6
Southern Ocean Medical Center	2	978	2.0	2.2	5.2 *	0.0 - 12.8
St. Barnabas Medical Center	64	6,033	10.6	3.0	20.4 **	17.8 - 23.0
St. Clare's Hospital-Denville	2	1,849	1.1	2.1	3.0 *	0.0 - 8.7
St. Clare's Hospital-Dover	4	640	6.3	2.2	16.0	6.6 - 25.3
St. Clare's Hospital-Sussex	0	39	0.0	2.3	0.0	0.0 - 37.6
St. Francis Medical Center-Trenton	8	1,251	6.4	2.2	17.0	10.2 - 23.9
St. Joseph's Hospital and Medical Center	32	4,805	6.7	2.2	17.5	14.1 - 21.0
St. Joseph's Wayne Hospital	5	945	5.3	2.2	13.6	5.9 - 21.4
St. Luke Warren Hospital	5	770	6.5	2.3	16.1	7.7 - 24.5
St. Mary's Hospital (Passaic)	17	1,822	9.3	2.4	22.1 **	16.7 - 27.4
St. Michael's Medical Center	9	1,892	4.8	2.6	10.5	5.4 - 15.5
St. Peter's University Hospital	17	2,714	6.3	2.5	14.2	9.9 - 18.5
Trinitas Regional Medical Center	5	1,968	2.5	2.3	6.4 *	1.1 - 11.7
UMDNJ-University Hospital	35	3,808	9.2	2.9	18.0	14.6 - 21.3
Underwood-Memorial Hospital	3	1,807	1.7	2.5	3.8 *	0.0 - 9.2
University Medical Center at Princeton	7	2,889	2.4	1.8	7.8 *	2.9 - 12.8
Valley Hospital	28	6,601	4.2	2.2	10.9 *	8.0 - 13.8
Virtua-Memorial Hospital Burlington Cty.	18	4,150	4.3	2.2	11.4	7.7 - 15.1
Virtua-West Jersey Hospital Berlin	0	254	0.0	2.3	0.0 *	0.0 - 14.6
Virtua-West Jersey Hospital Marlton	7	3,550	2.0	1.8	6.4 *	1.9 - 10.9
Virtua-West Jersey Hospital Voorhees	17	3,031	5.6	2.4	13.4	9.3 - 17.6

Source: New Jersey 2012 UB Data.

*: Statistically significantly below statewide average (i.e., better than statewide average).

**: Statistically significantly above statewide average (i.e., worse than statewide average).

Δ : Expected rate is the rate the hospital would have if it had the same case-mix (e.g., age, gender, DRG, and comorbidity categories) as the reference or statewide population. If the observed rate is higher than the expected rate (i.e., the ratio of observed to expected is greater than 1.0), it suggests that the hospital performed worse than the reference population on that indicator.

PSI.12- Perioperative pulmonary embolism (PE) or deep vein thrombosis (DVT)

- This indicator measures incidences of perioperative pulmonary embolism (blood clot in the lungs) or deep vein thrombosis (blood clot in a large vein) occurring during a surgical procedure. The rate, which is risk-adjusted for age, sex, DRG, and comorbidity, is defined as the number of perioperative pulmonary embolism (PE) or deep vein thrombosis (DVT) cases per 1,000 surgical discharges.
- The numerator includes discharges, among cases meeting the inclusion and exclusion rules for the denominator, with a secondary ICD-9-CM diagnosis codes for deep vein thrombosis or pulmonary embolism.
- The denominator includes all surgical discharges age 18 and older with ICD-9-CM procedure codes for an operating room procedure, defined by specific DRG or MS-DRG codes.
- The following cases are excluded from the denominator or from rate calculation:
 - cases with secondary diagnosis for pulmonary embolism or deep vein thrombosis, present on admission (pre-existing conditions);
 - cases with principal diagnosis for pulmonary embolism or deep vein thrombosis;
 - cases in which interruption of vena cava is the only operating room procedure or in which interruption of vena cava occurs before or on the same day as the first operating room procedure (*if day of procedure is not available in the input data file, the rate may be slightly lower than if the information was available*); and
 - obstetric discharges (MDC 14 - pregnancy, childbirth and the puerperium).
- Table 5 presents the number of perioperative pulmonary embolism or deep vein thrombosis cases among all surgical discharges age 18 and older by hospital, observed rates, expected rates, risk-adjusted rates, and the 95% confidence intervals computed for the risk-adjusted rates. Statewide, there were 1,315 cases of perioperative pulmonary embolism or deep vein thrombosis reported in 2012 for a statewide risk-adjusted rate of 5.5 per 1,000 surgical discharges.

Table 5. Perioperative Pulmonary Embolism or Deep Vein Thrombosis (per 1,000 surgical discharges)

Hospital	# of cases	# of discharges	Obs. rate	Exp. Rate Δ	Risk-adjusted rate	95% Confidence interval
Statewide	1,315	191,026	6.9	5.5	5.5	5.3 - 5.8
Atlanticare Regional Medical Center-City	12	1,878	6.4	7.1	3.9	1.7 - 6.2
Atlanticare Regional Medical Center-Mainland	28	4,817	5.8	6.8	3.7	2.3 - 5.2
Bayonne Medical Center	1	783	1.3	5.3	1.0 *	0.0 - 5.1
Bayshore Community Hospital	3	1,097	2.7	5.7	2.1	0.0 - 5.5
Bergen Regional Medical Center	6	169	35.5	3.6	42.9 **	32.0 - 53.8
Cape Regional Medical Center	5	1,197	4.2	8.7	2.1 *	0.0 - 4.6
Capital Health Medical Center - Hopewell	19	2,268	8.4	5.5	6.6	4.3 - 9.0
Capital Health Regional Medical Center	53	1,654	32.0	11.7	12.0 **	10.2 - 13.9
CentraState Medical Center	9	2,644	3.4	4.8	3.1 *	0.7 - 5.4
Chilton Memorial Hospital	8	1,865	4.3	6.5	2.9 *	0.6 - 5.2
Christ Hospital	8	1,161	6.9	3.7	8.1	4.0 - 12.1
Clara Maass Medical Center	11	3,612	3.0	4.2	3.1 *	1.0 - 5.3
Community Medical Center	21	4,494	4.7	5.8	3.5 *	1.9 - 5.2
Cooper Hospital/University Medical Center	79	6,802	11.6	5.4	9.4 **	8.0 - 10.8
Deborah Heart and Lung Center	9	2,784	3.2	4.2	3.4	0.9 - 5.9
East Orange General Hospital	2	539	3.7	4.6	3.6	0.0 - 9.0
Englewood Hospital and Medical Center	10	5,085	2.0	4.0	2.2 *	0.3 - 4.1
Hackensack University Medical Center	93	10,318	9.0	6.9	5.7	4.8 - 6.7
Hackettstown Regional Medical Center	5	799	6.3	5.3	5.2	1.2 - 9.2
Hoboken University Medical Center	6	859	7.0	3.8	8.1	3.4 - 12.7
Holy Name Medical Center	21	2,923	7.2	5.3	5.9	3.8 - 8.1
Hunterdon Medical Center	7	1,369	5.1	4.4	5.1	1.6 - 8.5
Jersey City Medical Center	7	2,091	3.3	5.0	2.9	0.4 - 5.5
Jersey Shore University Medical Center	33	6,535	5.0	7.0	3.1 *	1.9 - 4.4
JFK Medical Center/Anthony M. Yelencis	71	3,603	19.7	5.0	17.4 **	15.4 - 19.4
Kennedy University Hospital - Cherry Hill	5	538	9.3	5.1	8.0	2.9 - 13.1
Kennedy University Hospital - Stratford	3	1,144	2.6	4.6	2.5	0.0 - 6.0
Kennedy University Hospital - Wash. Twp.	29	2,603	11.1	6.5	7.5	5.5 - 9.5
Kimball Medical Center	5	778	6.4	5.9	4.8	0.9 - 8.7
Lourdes Medical Center of Burlington Cty.	3	1,192	2.5	4.4	2.5	0.0 - 6.1
Meadowlands Hospital Medical Center	1	481	2.1	3.3	2.8	0.0 - 9.5
Memorial Hospital of Salem County	0	497	0.0	2.7	0.0	0.0 - 7.4
Monmouth Medical Center	14	2,685	5.2	4.7	4.8	2.5 - 7.2
Morristown Memorial Hospital	83	11,399	7.3	5.4	5.9	4.8 - 6.9
Mountainside Hospital	14	1,850	7.6	4.8	6.9	4.1 - 9.7
Newark Beth Israel Medical Center	39	3,640	10.7	6.2	7.6 **	5.8 - 9.3
Newton Medical Center	3	982	3.1	5.3	2.5	0.0 - 6.2
Ocean Medical Center - Bricktown	15	3,020	5.0	6.0	3.6	1.7 - 5.6
Our Lady of Lourdes Medical Center	13	3,840	3.4	4.2	3.6	1.5 - 5.6
Overlook Medical Center	86	5,426	15.8	6.3	10.9 **	9.5 - 12.4
Palisades Medical Center - NY PHS	1	1,088	0.9	3.7	1.1 *	0.0 - 5.3
Raritan Bay Medical Center-Old Bridge	5	706	7.1	5.1	6.1	1.7 - 10.4
Raritan Bay Medical Center-Perth Amboy	6	1,112	5.4	4.6	5.1	1.5 - 8.7

Table 5. Perioperative Pulmonary Embolism or Deep Vein Thrombosis (per 1,000 surgical discharges)

Hospital	# of cases	# of discharges	Obs. rate	Exp. Rate Δ	Risk-adjusted rate	95% Confidence interval
Statewide	1,315	191,026	6.9	5.5	5.5	5.3 - 5.8
Riverview Medical Center	11	2,899	3.8	7.5	2.2 *	0.5 - 4.0
RWJ University Hospital	95	8,839	10.7	5.7	8.2 **	7.1 - 9.4
RWJ University Hospital at Hamilton	7	2,213	3.2	5.9	2.4 *	0.1 - 4.7
RWJ University Hospital at Rahway	2	942	2.1	6.7	1.4 *	0.0 - 4.7
Shore Medical Center	15	2,110	7.1	5.0	6.2	3.6 - 8.7
Somerset Medical Center	26	2,832	9.2	4.6	8.6 **	6.3 - 11.0
South Jersey Healthcare Regional MC	7	2,751	2.5	5.0	2.2 *	0.0 - 4.4
South Jersey Hospital-Elmer	0	740	0.0	5.3	0.0 *	0.0 - 4.3
Southern Ocean Medical Center	4	996	4.0	7.2	2.4	0.0 - 5.5
St. Barnabas Medical Center	55	6,217	8.8	5.3	7.3 **	5.8 - 8.7
St. Clare's Hospital-Denville	9	1,910	4.7	4.8	4.3	1.5 - 7.1
St. Clare's Hospital-Dover	1	649	1.5	6.1	1.1 *	0.0 - 5.1
St. Clare's Hospital-Sussex	0	39	0.0	3.6	0.0	0.0 - 22.8
St. Francis Medical Center-Trenton	8	1,275	6.3	3.8	7.1	3.3 - 10.9
St. Joseph's Hospital and Medical Center	8	4,970	1.6	3.9	1.8 *	0.0 - 3.8
St. Joseph's Wayne Hospital	1	978	1.0	6.1	0.7 *	0.0 4.2
St. Luke Warren Hospital	6	787	7.6	5.3	6.3	2.2 - 10.5
St. Mary's Hospital (Passaic)	8	1,866	4.3	4.0	4.6	1.5 - 7.7
St. Michael's Medical Center	6	1,936	3.1	5.0	2.7 *	0.0 - 5.4
St. Peter's University Hospital	21	2,807	7.5	4.6	7.1	4.8 - 9.5
Trinitas Regional Medical Center	9	2,044	4.4	4.3	4.5	1.6 - 7.3
UMDNJ-University Hospital	59	4,022	14.7	5.9	10.8 **	9.1 - 12.6
Underwood-Memorial Hospital	8	1,859	4.3	4.8	4.0	1.1 - 6.8
University Medical Center at Princeton	7	2,979	2.3	4.5	2.3 *	0.0 - 4.6
Valley Hospital	54	6,802	7.9	5.2	6.7	5.2 - 8.1
Virtua-Memorial Hospital Burlington Cty.	16	4,249	3.8	4.8	3.4 *	1.6 - 5.3
Virtua-West Jersey Hospital Berlin	0	267	0.0	6.6	0.0	0.0 - 6.2
Virtua-West Jersey Hospital Marlton	10	3,622	2.8	5.3	2.3 *	0.4 - 4.1
Virtua-West Jersey Hospital Voorhees	10	3,099	3.2	5.3	2.7 *	0.6 - 4.7

Source: New Jersey 2012 UB Data.

*: Statistically significantly below statewide average (i.e., better than statewide average).

** : Statistically significantly above statewide average (i.e., worse than statewide average).

Δ : Expected rate is the rate the hospital would have if it had the same case-mix (e.g., age, gender, DRG, and comorbidity categories) as the reference or statewide population. If the observed rate is higher than the expected rate (i.e., the ratio of observed to expected is greater than 1.0), it suggests that the hospital performed worse than the reference population on that indicator.

PSI.13 – Postoperative sepsis

- This indicator flags how often hospitalized patients get a serious bloodstream infection (nosocomial postoperative sepsis). A serious infection of the bloodstream caused by toxin-producing bacteria, known as sepsis, can occur after surgery. The rate, which is risk-adjusted for age, sex, DRG, and comorbidity categories, is defined as the number of sepsis cases per 1,000 elective surgery patients with an operating room procedure and a length of stay of 4 days or more.
- The numerator includes discharges with secondary ICD-9-CM diagnosis codes for sepsis while the denominator includes all elective surgical discharges age 18 and older defined by specific DRG or MS-DRG codes with admission type recorded as elective.
- The following cases are excluded from the denominator or from rate calculation:
 - cases with a principal ICD-9-CM diagnosis code (or secondary diagnosis present on admission) for sepsis;
 - cases with a principal ICD-9-CM diagnosis code (or secondary diagnosis present on admission) for infection;
 - cases with any listed ICD-9-CM diagnosis codes or any listed ICD-9-CM procedure codes for immunocompromised state;
 - cases with any listed ICD-9-CM diagnosis codes for cancer;
 - MDC 14 (pregnancy, childbirth, and puerperium); and
 - cases with a length of stay of less than 4 days.
- Table 6 shows the number of postoperative sepsis cases among elective surgery patients by hospital, as well as the observed, expected and risk-adjusted rates along with their corresponding 95% confidence intervals. Statewide, there were 157 postoperative sepsis cases reported in 2012 for a statewide risk-adjusted rate of 11.1 per 1,000 elective discharges.

Table 6. Postoperative Sepsis (per 1,000 elective surgical discharges)

Hospital	# of cases	# of discharges	Obs. rate	Exp. Rate Δ	Risk-adjusted rate	95% Confidence interval
Statewide	157	13,752	11.4	12.2	11.1	9.3 - 12.8
Atlanticare Regional Medical Center-City	0	70	0.0	11.2	0.0	0.0 - 25.9
Atlanticare Regional Medical Center-Mainland	1	394	2.5	10.1	3.0	0.0 - 14.3
Bayonne Medical Center	1	35	28.6	13.0	26.0	0.0 - 60.0
Bayshore Community Hospital	0	10	0.0	16.7	0.0 [^]	0.0 - 55.1
Bergen Regional Medical Center	0	50	0.0	6.1	0.0	0.0 - 41.5
Cape Regional Medical Center	0	38	0.0	10.8	0.0	0.0 - 35.8
Capital Health Medical Center - Hopewell	0	171	0.0	8.8	0.0	0.0 - 18.6
Capital Health Regional Medical Center	0	40	0.0	7.0	0.0	0.0 - 43.5
CentraState Medical Center	3	88	34.1	11.5	35.0 ^{**}	12.3 - 57.8
Chilton Memorial Hospital	0	94	0.0	10.3	0.0	0.0 - 23.2
Christ Hospital	3	51	58.8	13.6	51.2 ^{**}	24.3 - 78.1
Clara Maass Medical Center	0	142	0.0	11.6	0.0	0.0 - 17.8
Community Medical Center	1	222	4.5	14.5	3.7	0.0 - 16.1
Cooper Hospital/University Medical Center	17	794	21.4	14.8	17.0	10.4 - 23.6
Deborah Heart and Lung Center	0	19	0.0	17.7	0.0 [^]	0.0 - 39.0
East Orange General Hospital	1	30	33.3	4.3	91.3 ^{**}	27.3 - 155.3
Englewood Hospital and Medical Center	4	590	6.8	11.1	7.2	0.0 - 16.2
Hackensack University Medical Center	7	823	8.5	12.0	8.4	1.1 - 15.7
Hackettstown Regional Medical Center	3	53	56.6	6.6	101.2 ^{**}	62.3 - 140.1
Hoboken University Medical Center	0	53	0.0	6.1	0.0	0.0 - 40.3
Holy Name Medical Center	3	169	17.8	8.4	24.9	5.7 - 44.1
Hunterdon Medical Center	0	60	0.0	13.5	0.0	0.0 - 25.3
Jersey City Medical Center	3	148	20.3	10.4	22.9	4.5 - 41.4
Jersey Shore University Medical Center	0	77	0.0	18.4	0.0	0.0 - 19.0
JFK Medical Center/Anthony M. Yelensics	5	367	13.6	10.7	15.0	3.5 - 26.5
Kennedy University Hospital - Cherry Hill	0	18	0.0	8.1	0.0 [^]	0.0 - 59.8
Kennedy University Hospital - Stratford	0	61	0.0	10.4	0.0	0.0 - 28.9
Kennedy University Hospital - Wash. Twp.	1	127	7.9	9.6	9.7	0.0 - 30.3
Kimball Medical Center	1	20	50.0	6.6	89.7 ^{^***}	26.3 - 153.1
Lourdes Medical Center of Burlington Cty.	0	54	0.0	13.4	0.0	0.0 - 26.8
Meadowlands Hospital Medical Center	0	9	0.0	8.0	0.0 [^]	0.0 - 85.9
Memorial Hospital of Salem County	1	29	34.5	8.0	51.1 [^]	3.2 - 99.0
Monmouth Medical Center	2	231	8.7	10.3	9.9	0.0 - 24.8
Morristown Memmorial Hospital	5	1,235	4.0	15.0	3.2 [*]	0.0 - 8.4
Mountainside Hospital	4	112	35.7	11.5	36.6 ^{**}	16.4 - 56.7
Newark Beth Israel Medical Center	4	309	12.9	13.6	11.3	0.1 - 22.4
Newton Medical Center	0	82	0.0	7.8	0.0	0.0 - 28.6
Ocean Medical Center - Bricktown	0	39	0.0	11.1	0.0	0.0 - 34.8
Our Lady of Lourdes Medical Center	9	653	13.8	16.7	9.8	2.9 - 16.6
Overlook Medical Center	4	350	11.4	10.5	12.9	0.9 - 24.9
Palisades Medical Center - NY PHS	1	58	17.2	13.2	15.4	0.0 - 41.4
Raritan Bay Medical Center-Old Bridge	0	15	0.0	18.2	0.0 [^]	0.0 - 43.4
Raritan Bay Medical Center-Perth Amboy	1	61	16.4	8.7	22.2	0.0 - 53.6
Riverview Medical Center	1	22	45.5	13.7	39.2 [^]	0.0 - 80.8

Table 6. Postoperative Sepsis (per 1,000 elective surgical discharges)

Hospital	# of cases	# of discharges	Obs. rate	Exp. Rate Δ	Risk-adjusted rate	95% Confidence interval
Statewide	157	13,752	11.4	12.2	11.1	9.3 - 12.8
RWJ University Hospital	10	885	11.3	11.6	11.5	4.4 - 18.7
RWJ University Hospital at Hamilton	1	151	6.6	9.4	8.3	0.0 - 27.5
RWJ University Hospital at Rahway	1	47	21.3	28.7	8.8	0.0 - 27.7
Shore Medical Center	0	120	0.0	9.3	0.0	0.0 - 21.7
Somerset Medical Center	4	250	16.0	12.4	15.2	2.3 - 28.2
South Jersey Healthcare Regional MC	1	218	4.6	13.7	4.0	0.0 - 17.1
South Jersey Hospital-Elmer	0	54	0.0	13.2	0.0	0.0 - 27.0
Southern Ocean Medical Center	0	2	0.0	.	.	- .
St. Barnabas Medical Center	5	554	9.0	12.9	8.2	0.0 - 16.7
St. Clare's Hospital-Denville	0	71	0.0	10.1	0.0	0.0 - 27.1
St. Clare's Hospital-Dover	0	14	0.0	17.0	0.0 ^	0.0 - 46.6
St. Clare's Hospital-Sussex	- .
St. Francis Medical Center-Trenton	5	130	38.5	12.0	37.7 **	19.4 55.9
St. Joseph's Hospital and Medical Center	9	435	20.7	10.5	23.3 **	12.6 - 34.1
St. Joseph's Wayne Hospital	1	73	13.7	9.3	17.4	0.0 - 45.2
St. Luke Warren Hospital	4	62	64.5	10.7	71.2 **	43.2 - 99.3
St. Mary's Hospital (Passaic)	2	153	13.1	15.8	9.8	0.0 - 24.2
St. Michael's Medical Center	5	173	28.9	14.5	23.6	9.1 - 38.0
St. Peter's University Hospital	3	191	15.7	12.3	15.1	0.2 - 30.0
Trinitas Regional Medical Center	0	93	0.0	11.0	0.0	0.0 - 22.7
UMDNJ-University Hospital	5	422	11.8	8.7	16.1	4.1 - 28.0
Underwood-Memorial Hospital	0	110	0.0	9.1	0.0	0.0 - 22.8
University Medical Center at Princeton	5	232	21.6	10.3	24.8	10.0 - 39.6
Valley Hospital	4	463	8.6	11.6	8.8	0.0 18.7
Virtua-Memorial Hospital Burlington Cty.	2	293	6.8	12.3	6.5	0.0 - 18.5
Virtua-West Jersey Hospital Berlin	0	4	0.0	10.4	0.0 ^	0.0 112.8
Virtua-West Jersey Hospital Marlton	2	333	6.0	9.7	7.3	0.0 - 19.9
Virtua-West Jersey Hospital Voorhees	2	146	13.7	15.4	10.5	0.0 - 25.5

Source: New Jersey 2012 UB Data.

*: Statistically significantly below statewide average (i.e., better than statewide average).

**: Statistically significantly above statewide average (i.e., worse than statewide average).

^: Rate is based on a denominator less than 30, and should be taken with caution.

Δ : Expected rate is the rate the hospital would have if it had the same case-mix (e.g., age, gender, DRG, and comorbidity categories) as the reference or statewide population. If the observed rate is higher than the expected rate (i.e., the ratio of observed to expected is greater than 1.0), it suggests that the hospital performed worse than the reference population on that indicator.

PSI.14 – Postoperative wound dehiscence

- This indicator flags cases of wound dehiscence (i.e. when surgical wound in the stomach or pelvic area is split open after an operation) in patients who have undergone abdominal and pelvic surgery. Some or all of these complications may require treatment with another major operation to fix the wound. Wound dehiscence following surgery is a medical error that can be avoided. The rate, which is risk-adjusted for age, sex, DRG, and comorbidity categories, is defined as the number of cases of re-closure of postoperative disruption of abdominal wall per 1,000 cases of abdominopelvic surgery.
- The numerator includes discharges with ICD-9-CM code (5461) for re-closure of postoperative disruption of abdominal wall in any procedure field, while the denominator includes all patients age 18 and older with any listed ICD-9-CM procedure codes for abdominopelvic surgery.
- The following cases are excluded from the denominator or from rate calculation:
 - cases where a procedure for re-closure of postoperative disruption of abdominal wall occurs before or on the same day as the first abdominopelvic surgery procedure (*if day of procedure is not available in the input data file, the rate may be slightly lower than if the information was available*);
 - with any-listed ICD-9-CM diagnosis codes or any-listed ICD-9-CM procedure codes for immunocompromised state;
 - cases where length of stay is less than 2 days; and
 - MDC 14 (pregnancy, childbirth, and puerperium).
- Table 7 shows the number of post-operative wound dehiscence cases among patients who have undergone abdominal and pelvic surgery by hospital, observed rates, expected rates, and risk-adjusted rates along with their corresponding 95% confidence intervals. Statewide, there were 62 postoperative wound dehiscence cases reported in 2012 for a statewide risk-adjusted rate of 1.7 per 1,000 abdominopelvic surgical discharges.

Table 7. Postoperative wound dehiscence (per 1,000 abdominopelvic surgical discharges)

Hospital	# of cases	# of discharges	Obs. rate	Exp. Rate Δ	Risk-adjusted rate	95% Confidence interval
Statewide	62	30,184	2.1	2.3	1.7	1.2 - 2.1
Atlanticare Regional Medical Center-City	2	307	6.5	2.1	5.9	1.3 - 10.5
Atlanticare Regional Medical Center-Mainland	0	343	0.0	2.5	0.0	0.0 - 4.0
Bayonne Medical Center	0	140	0.0	2.9	0.0	0.0 - 5.8
Bayshore Community Hospital	0	191	0.0	2.9	0.0	0.0 - 4.9
Bergen Regional Medical Center	0	16	0.0	3.2	0.0 ^	0.0 - 16.0
Cape Regional Medical Center	1	250	4.0	2.7	2.7	0.0 - 7.2
Capital Health Medical Center - Hopewell	1	498	2.0	2.0	1.9	0.0 - 5.6
Capital Health Regional Medical Center	1	190	5.3	2.6	3.8	0.0 - 9.1
CentraState Medical Center	0	496	0.0	2.6	0.0	0.0 - 3.2
Chilton Memorial Hospital	1	316	3.2	2.5	2.4	0.0 - 6.5
Christ Hospital	1	230	4.3	1.9	4.2	0.0 - 9.7
Clara Maass Medical Center	0	581	0.0	2.6	0.0	0.0 - 3.0
Community Medical Center	3	887	3.4	2.9	2.2	0.0 - 4.5
Cooper Hospital/University Medical Center	6	1,401	4.3	1.9	4.2 **	2.0 - 6.5
Deborah Heart and Lung Center	0	16	0.0	1.9	0.0 ^	0.0 - 20.8
East Orange General Hospital	0	106	0.0	3.4	0.0	0.0 - 6.1
Englewood Hospital and Medical Center	0	630	0.0	2.0	0.0	0.0 - 3.3
Hackensack University Medical Center	0	1,293	0.0	2.4	0.0	0.0 - 2.1
Hackettstown Regional Medical Center	0	114	0.0	3.0	0.0	0.0 - 6.3
Hoboken University Medical Center	0	111	0.0	2.8	0.0	0.0 - 6.5
Holy Name Medical Center	1	575	1.7	1.9	1.7	0.0 - 5.3
Hunterdon Medical Center	0	238	0.0	2.3	0.0	0.0 - 5.0
Jersey City Medical Center	0	280	0.0	2.6	0.0	0.0 - 4.3
Jersey Shore University Medical Center	0	723	0.0	3.7	0.0	0.0 - 2.2
JFK Medical Center/Anthony M. Yelensics	0	699	0.0	2.1	0.0	0.0 - 3.0
Kennedy University Hospital - Cherry Hill	1	69	14.5	2.5	10.8 **	2.0 - 19.6
Kennedy University Hospital - Stratford	1	208	4.8	1.8	4.9	0.0 - 10.8
Kennedy University Hospital - Wash. Twp.	2	358	5.6	2.4	4.4	0.4 - 8.4
Kimball Medical Center	0	174	0.0	2.1	0.0	0.0 - 6.0
Lourdes Medical Center of Burlington Cty.	0	231	0.0	2.6	0.0	0.0 - 4.7
Meadowlands Hospital Medical Center	0	113	0.0	1.2	0.0	0.0 - 9.9
Memorial Hospital of Salem County	0	130	0.0	1.5	0.0	0.0 - 8.2
Monmouth Medical Center	1	581	1.7	2.4	1.4	0.0 - 4.5
Morristown Memmorial Hospital	0	1,169	0.0	2.5	0.0	0.0 - 2.1
Mountainside Hospital	0	394	0.0	2.2	0.0	0.0 - 3.9
Newark Beth Israel Medical Center	0	510	0.0	1.5	0.0	0.0 - 4.2
Newton Medical Center	2	188	10.6	2.9	6.9 **	1.9 - 11.8
Ocean Medical Center - Bricktown	0	494	0.0	2.9	0.0	0.0 - 3.0
Our Lady of Lourdes Medical Center	1	466	2.1	2.3	1.8	0.0 - 5.3
Overlook Medical Center	3	1,037	2.9	2.1	2.5	0.1 - 5.0
Palisades Medical Center - NY PHS	1	194	5.2	2.5	3.9	0.0 - 9.2
Raritan Bay Medical Center-Old Bridge	0	106	0.0	2.8	0.0	0.0 - 6.7
Raritan Bay Medical Center-Perth Amboy	0	193	0.0	1.6	0.0	0.0 - 6.5
Riverview Medical Center	2	502	4.0	2.9	2.6	0.0 - 5.6

Table 7. Postoperative wound dehiscence (per 1,000 abdominopelvic surgical discharges)

Hospital	# of cases	# of discharges	Obs. rate	Exp. Rate Δ	Risk-adjusted rate	95% Confidence interval
Statewide	62	30,184	2.1	2.3	1.7	1.2 - 2.1
RWJ University Hospital	1	1,243	0.8	2.1	0.7	0.0 - 3.0
RWJ University Hospital at Hamilton	0	428	0.0	2.4	0.0	0.0 - 3.6
RWJ University Hospital at Rahway	1	151	6.6	3.0	4.2	0.0 - 9.7
Shore Medical Center	0	275	0.0	2.5	0.0	0.0 - 4.4
Somerset Medical Center	1	388	2.6	2.6	1.9	0.0 - 5.5
South Jersey Healthcare Regional MC	1	607	1.6	1.7	1.8	0.0 - 5.4
South Jersey Hospital-Elmer	0	58	0.0	2.5	0.0	0.0 - 9.6
Southern Ocean Medical Center	1	217	4.6	2.8	3.1	0.0 - 7.8
St. Barnabas Medical Center	6	1,225	4.9	1.7	5.3 **	2.8 - 7.9
St. Clare's Hospital-Denville	0	278	0.0	2.2	0.0	0.0 - 4.6
St. Clare's Hospital-Dover	0	134	0.0	2.1	0.0	0.0 - 6.8
St. Clare's Hospital-Sussex	0	2	0.0	.	.	. - .
St. Francis Medical Center-Trenton	1	168	6.0	2.4	4.6	0.0 - 10.4
St. Joseph's Hospital and Medical Center	3	633	4.7	2.1	4.3	1.1 - 7.5
St. Joseph's Wayne Hospital	0	224	0.0	2.9	0.0	0.0 - 4.5
St. Luke Warren Hospital	0	190	0.0	2.4	0.0	0.0 - 5.4
St. Mary's Hospital (Passaic)	1	297	3.4	2.6	2.4	0.0 - 6.6
St. Michael's Medical Center	0	289	0.0	2.2	0.0	0.0 - 4.6
St. Peter's University Hospital	4	809	4.9	1.7	5.6 **	2.4 - 8.7
Trinitas Regional Medical Center	2	383	5.2	1.9	5.0	0.8 - 9.3
UMDNJ-University Hospital	2	515	3.9	2.3	3.2	0.0 - 6.6
Underwood-Memorial Hospital	3	313	9.6	2.5	7.2 **	3.0 - 11.3
University Medical Center at Princeton	0	371	0.0	2.2	0.0	0.0 - 4.0
Valley Hospital	2	1,075	1.9	2.4	1.4	0.0 - 3.7
Virtua-Memorial Hospital Burlington Cty.	0	859	0.0	2.3	0.0	0.0 - 2.6
Virtua-West Jersey Hospital Berlin	0	42	0.0	4.1	0.0	0.0 - 8.8
Virtua-West Jersey Hospital Marlton	2	370	5.4	2.6	3.9	0.2 - 7.6
Virtua-West Jersey Hospital Voorhees	0	892	0.0	2.0	0.0	0.0 - 2.7

Source: New Jersey 2012 UB Data.

** : Statistically significantly above statewide average (i.e., worse than statewide average).

^ : Rates based on denominators less than 30 should be taken with caution.

Δ : Expected rate is the rate the hospital would have if it had the same case-mix (e.g., age, gender, DRG, and comorbidity categories) as the reference or statewide population. If the observed rate is higher than the expected rate (i.e., the ratio of observed to expected is greater than 1.0), it suggests that the hospital performed worse than the reference population on that indicator.

PSI.15 - Accidental puncture or laceration

- This indicator measures the occurrence of complications that arise due to technical difficulties in medical care, specifically, those involving an accidental puncture or laceration. The rate, which is risk-adjusted for age, sex, DRG, and comorbidity categories, is defined as the number of patients who had an accidental cut or lacerations (secondary diagnosis) during a medical procedure per 1,000 discharges.
- The numerator refers to all discharges with any secondary ICD-9-CM code denoting technical difficulty (e.g., accidental cut, puncture, perforation, or laceration) during a procedure.
- The denominator refers to all medical and surgical discharges age 18 and older defined by specific DRG or MS-DRG codes, excluding cases:
 - with a principal ICD-9-CM diagnosis code (or secondary diagnosis present on admission) for accidental puncture or laceration during a procedure;
 - with any listed ICD-9-CM procedure codes for spine surgery; and
 - MDC 14 (pregnancy, childbirth, and puerperium).
- Table 8 shows the number of cases of accidental puncture or laceration among all discharges with ICD-9-CM code denoting technical difficulty (e.g., accidental cut, puncture, perforation, or laceration) in any secondary diagnosis field by hospital along with observed and expected rates as well as risk-adjusted rates with their corresponding 95% confidence intervals. Statewide, there were 969 cases of accidental punctures or lacerations reported in 2012 for a risk-adjusted rate of 1.2 per 1,000 discharges.

Table 8. Accidental puncture or laceration (per 1,000 medical and surgical discharges)

Hospital	# of cases	# of discharges	Obs. rate	Exp. Rate Δ	Risk-adjusted rate	95% Confidence interval
Statewide	969	732,958	1.3	2.6	1.2	1.1 - 1.3
Atlanticare Regional Medical Center-City	4	9,167	0.4	1.8	0.6	0.0 - 1.7
Atlanticare Regional Medical Center-Mainland	7	15,414	0.5	2.0	0.5	0.0 - 1.4
Bayonne Medical Center	1	5,138	0.2	2.0	0.2	0.0 - 1.7
Bayshore Community Hospital	3	6,912	0.4	1.7	0.6	0.0 - 2.0
Bergen Regional Medical Center	0	6,147	0.0	0.4	0.0	0.0 - 3.1
Cape Regional Medical Center	7	8,292	0.8	1.5	1.4	0.0 - 2.7
Capital Health Medical Center - Hopewell	26	7,489	3.5	3.5	2.4 **	1.5 - 3.3
Capital Health Regional Medical Center	2	9,619	0.2	1.3	0.4	0.0 - 1.7
CentraState Medical Center	26	11,371	2.3	2.6	2.2 **	1.3 - 3.0
Chilton Memorial Hospital	21	8,359	2.5	2.3	2.7 **	1.6 - 3.7
Christ Hospital	9	7,144	1.3	1.8	1.7	0.4 - 3.0
Clara Maass Medical Center	10	13,992	0.7	2.4	0.7	0.0 - 1.5
Community Medical Center	24	22,902	1.0	2.3	1.1	0.5 - 1.8
Cooper Hospital/University Medical Center	48	18,852	2.5	3.9	1.6	1.0 - 2.1
Deborah Heart and Lung Center	2	4,512	0.4	4.8	0.2	0.0 - 1.2
East Orange General Hospital	0	7,170	0.0	1.1	0.0	0.0 - 1.7
Englewood Hospital and Medical Center	6	12,698	0.5	3.6	0.3 *	0.0 - 1.0
Hackensack University Medical Center	60	27,501	2.2	3.3	1.6	1.1 - 2.1
Hackettstown Regional Medical Center	2	3,694	0.5	2.0	0.7	0.0 - 2.4
Hoboken University Medical Center	4	4,419	0.9	1.6	1.3	0.0 - 3.1
Holy Name Medical Center	14	11,252	1.2	2.8	1.1	0.2 - 1.9
Hunterdon Medical Center	2	6,251	0.3	2.1	0.4	0.0 - 1.7
Jersey City Medical Center	6	11,383	0.5	1.9	0.7	0.0 - 1.7
Jersey Shore University Medical Center	49	18,766	2.6	3.8	1.7	1.1 - 2.2
JFK Medical Center/Anthony M. Yelencsics	7	13,650	0.5	2.8	0.5	0.0 - 1.2
Kennedy University Hospital - Cherry Hill	1	7,100	0.1	0.9	0.4	0.0 - 2.3
Kennedy University Hospital - Stratford	5	6,840	0.7	1.8	1.0	0.0 - 2.4
Kennedy University Hospital - Wash. Twp.	11	12,663	0.9	1.9	1.1	0.2 - 2.1
Kimball Medical Center	6	8,635	0.7	1.1	1.5	0.0 - 3.0
Lourdes Medical Center of Burlington Cty.	8	6,638	1.2	2.0	1.5	0.2 - 2.8
Meadowlands Hospital Medical Center	3	2,071	1.4	2.7	1.3	0.0 - 3.3
Memorial Hospital of Salem County	3	3,572	0.8	1.7	1.2	0.0 - 3.1
Monmouth Medical Center	9	8,971	1.0	3.2	0.8	0.0 - 1.6
Morristown Memorial Hospital	45	24,234	1.9	4.3	1.1	0.6 - 1.5
Mountainside Hospital	9	7,920	1.1	2.6	1.1	0.0 - 2.1
Newark Beth Israel Medical Center	16	13,800	1.2	2.9	1.0	0.2 - 1.7
Newton Medical Center	4	6,216	0.6	1.8	0.9	0.0 - 2.3
Ocean Medical Center - Bricktown	30	11,921	2.5	2.6	2.3 **	1.5 - 3.2
Our Lady of Lourdes Medical Center	6	11,359	0.5	3.4	0.4	0.0 - 1.1
Overlook Medical Center	54	16,344	3.3	3.6	2.2 **	1.6 - 2.8
Palisades Medical Center - NY PHS	8	7,650	1.0	1.6	1.6	0.2 - 2.9
Raritan Bay Medical Center-Old Bridge	2	4,593	0.4	1.5	0.7	0.0 - 2.5
Raritan Bay Medical Center-Perth Amboy	6	6,844	0.9	1.8	1.2	0.0 - 2.5
Riverview Medical Center	16	9,400	1.7	3.3	1.3	0.4 - 2.1

Table 8. Accidental puncture or laceration (per 1,000 medical and surgical discharges)

Hospital	# of cases	# of discharges	Obs. rate	Exp. Rate Δ	Risk-adjusted rate	95% Confidence interval
Statewide	969	732,958	1.3	2.6	1.2	1.1 - 1.3
RWJ University Hospital	57	25,693	2.2	3.6	1.5	1.0 - 2.0
RWJ University Hospital at Hamilton	30	9,913	3.0	2.2	3.3 **	2.3 - 4.3
RWJ University Hospital at Rahway	1	6,386	0.2	1.8	0.2	0.0 - 1.6
Shore Medical Center	9	8,345	1.1	2.1	1.2	0.1 - 2.3
Somerset Medical Center	16	12,269	1.3	2.5	1.3	0.4 - 2.1
South Jersey Healthcare Regional MC	20	12,665	1.6	2.3	1.7	0.8 - 2.6
South Jersey Hospital-Elmer	1	3,262	0.3	1.6	0.5	0.0 - 2.6
Southern Ocean Medical Center	7	5,437	1.3	2.1	1.5	0.1 - 2.9
St. Barnabas Medical Center	35	18,008	1.9	3.8	1.2	0.7 - 1.8
St. Clare's Hospital-Denville	6	9,594	0.6	2.2	0.7	0.0 - 1.7
St. Clare's Hospital-Dover	0	3,740	0.0	2.0	0.0	0.0 - 1.7
St. Clare's Hospital-Sussex	0	563	0.0	0.7	0.0	0.0 - 7.8
St. Francis Medical Center-Trenton	13	6,204	2.1	1.8	2.8 **	1.4 - 4.2
St. Joseph's Hospital and Medical Center	28	21,364	1.3	2.1	1.5	0.8 - 2.2
St. Joseph's Wayne Hospital	1	5,380	0.2	2.1	0.2	0.0 - 1.6
St. Luke Warren Hospital	6	4,490	1.3	2.1	1.5	0.0 - 3.1
St. Mary's Hospital (Passaic)	11	7,310	1.5	2.5	1.5	0.4 - 2.6
St. Michael's Medical Center	7	9,074	0.8	2.1	0.9	0.0 - 2.0
St. Peter's University Hospital	25	9,742	2.6	3.7	1.7	0.9 - 2.4
Trinitas Regional Medical Center	7	10,301	0.7	2.1	0.8	0.0 - 1.8
UMDNJ-University Hospital	12	12,352	1.0	2.5	0.9	0.1 - 1.8
Underwood-Memorial Hospital	4	9,989	0.4	2.1	0.5	0.0 - 1.5
University Medical Center at Princeton	16	9,880	1.6	2.8	1.4	0.5 - 2.3
Valley Hospital	17	21,584	0.8	3.2	0.6	0.0 - 1.2
Virtua-Memorial Hospital Burlington Cty.	18	13,661	1.3	3.0	1.1	0.3 - 1.8
Virtua-West Jersey Hospital Berlin	0	3,637	0.0	1.0	0.0	0.0 - 2.5
Virtua-West Jersey Hospital Marlton	12	10,139	1.2	2.7	1.1	0.2 - 2.0
Virtua-West Jersey Hospital Voorhees	28	13,111	2.1	3.2	1.6	0.9 - 2.3

Source: New Jersey 2012 UB Data.

*: Statistically significantly below statewide average (i.e., better than statewide average).

** : Statistically significantly above statewide average (i.e., worse than statewide average).

Δ : Expected rate is the rate the hospital would have if it had the same case-mix (e.g., age, gender, DRG, and comorbidity categories) as the reference or statewide population. If the observed rate is higher than the expected rate (i.e., the ratio of observed to expected is greater than 1.0), it suggests that the hospital performed worse than the reference population on that indicator.

PSI.16 - Transfusion reaction

- This indicator flags cases of major reactions due to transfusions (ABO and Rh). It measures the occurrence of major reactions to blood transfusions. Using the wrong type of blood or blood substitute are examples of why this type of medical error may occur. The indicator is measured using volume of occurrence – not a rate. It tells you the number of patients who had a transfusion reaction during hospital stay. It is considered a never-event and happens very rarely.
- The measure refers to discharges 18 years and older or in MDC 14 with ICD-9-CM codes for transfusion reaction in any secondary diagnosis field of all medical and surgical discharges defined by specific DRGs or MS-DRGs.
- Patients with principal diagnosis of transfusion reaction or secondary diagnosis present on admission are excluded from the measure.
- Transfusion reaction is considered a never-event and happens very rarely. Statewide, only 1 case of transfusion reaction was reported in 2012. It was reported by Hackensack University Medical Center.

PSI.17 - Birth trauma - injury to neonate

- This indicator flags cases of birth trauma among all newborns in a hospital. Birth trauma (injury to neonate) is caused by medical complications during labor and delivery. The rate is defined as the number of cases of birth trauma (injury to neonate) per 1,000 live births. No risk-adjustment is made to this indicator and no significance tests performed as a result.
- The numerator includes discharges with ICD-9-CM code for birth trauma in any diagnosis field excluding infants with a subdural or cerebral hemorrhage and any diagnosis code of pre-term infant (denoting birth weight of <2,500 grams and <37 gestation weeks or <=34 gestation weeks). It also excludes infants with injury to skeleton (CD-9-CM codes - 7673 and 7674) and infants with diagnosis code of osteogenesis imperfecta (CD-9-CM code - 75651).
- The denominator includes all livebirths (newborns).
- Table 9 shows the number of 'birth trauma - injury to neonate' cases among infants born alive and the observed rates by hospital. Statewide, there were 170 cases of 'birth trauma - injury to neonate' reported in 2012 for an observed rate of 1.7 per 1,000 live births.

Table 9. Birth Trauma - Injury to Neonate (per 1,000 livebirths)

(This indicator is calculated using the Pediatric Quality Indicators (PDIs) Module because it is based on pediatric discharges.)

Hospital	# of cases	# of livebirths	Observed rate
Statewide	170	98,181	1.7
Atlanticare Regional Medical Center-City	0	30	0.0
Atlanticare Regional Medical Center-Mainland	5	2,383	2.1
Cape Regional Medical Center	0	525	0.0
Capital Health Medical Center - Hopewell	3	1,896	1.6
Capital Health Regional Medical Center	0	351	0.0
CentraState Medical Center	1	1,490	0.7
Chilton Memorial Hospital	2	893	2.2
Christ Hospital	3	1,368	2.2
Clara Maass Medical Center	0	1,524	0.0
Community Medical Center	0	1,550	0.0
Cooper Hospital/University Medical Center	10	2,065	4.8
Englewood Hospital and Medical Center	2	1,936	1.0
Hackensack University Medical Center	16	5,928	2.7
Hackettstown Regional Medical Center	2	480	4.2
Hoboken University Medical Center	0	968	0.0
Holy Name Medical Center	1	1,485	0.7
Hunterdon Medical Center	1	947	1.1
Jersey City Medical Center	4	1,637	2.4
Jersey Shore University Medical Center	3	1,907	1.6
JFK Medical Center/Anthony M. Yelensics	8	2,634	3.0
Kennedy University Hospital - Wash. Twp.	2	1,018	2.0
Kimball Medical Center	1	894	1.1
Meadowlands Hospital Medical Center	1	839	1.2
Memorial Hospital of Salem County	0	173	0.0
Monmouth Medical Center	7	4,344	1.6
Morristown Memorial Hospital	9	4,066	2.2
Mountainside Hospital	11	1,256	8.8
Newark Beth Israel Medical Center	8	3,342	2.4
Newton Medical Center	2	540	3.7
Ocean Medical Center - Bricktown	2	948	2.1
Our Lady of Lourdes Medical Center	1	869	1.2
Overlook Medical Center	1	2,399	0.4
Palisades Medical Center - NY PHS	1	1,505	0.7

Table 9. Birth Trauma - Injury to Neonate (per 1,000 livebirths)

(This indicator is calculated using the Pediatric Quality Indicators (PDIs) Module because it is based on pediatric discharges.)

Hospital	# of cases	# of livebirths	Observed rate
Statewide	170	98,181	1.7
Raritan Bay Medical Center-Perth Amboy	0	1,248	0.0
Riverview Medical Center	0	1,351	0.0
RWJ University Hospital	4	2,110	1.9
RWJ University Hospital at Hamilton	2	1,076	1.9
Shore Medical Center	0	1,009	0.0
Somerset Medical Center	1	991	1.0
South Jersey Healthcare Regional MC	3	1,917	1.6
South Jersey Hospital-Elmer	1	279	3.6
Southern Ocean Medical Center	0	321	0.0
St. Barnabas Medical Center	12	5,411	2.2
St. Clare's Hospital-Denville	1	1,282	0.8
St. Joseph's Hospital and Medical Center	4	3,227	1.2
St. Mary's Hospital (Passaic)	0	875	0.0
St. Peter's University Hospital	9	5,590	1.6
Trinitas Regional Medical Center	0	2,137	0.0
UMDNJ-University Hospital	4	1,422	2.8
Underwood-Memorial Hospital	2	874	2.3
University Medical Center at Princeton	4	1,927	2.1
Valley Hospital	4	3,050	1.3
Virtua-Memorial Hospital Burlington Cty.	4	2,494	1.6
Virtua-West Jersey Hospital Voorhees	8	5,400	1.5

Source: New Jersey 2012 UB Data.

Note that this indicator is measured using observed rate only. Risk-adjustment is no more made on this indicator.

Newborn or livebirth is defined as any neonate with either 1) an ICD-9-CM diagnosis code for an in-hospital live-born birth or 2) an admission type of newborn (ATYPE=4), age in days at admission equaling zero, and not an ICD-9-CM diagnosis code for an out-of-hospital birth. A neonate is defined as any discharge with age in days at admission between zero and 28 days (inclusive). If age in days is missing, then a neonate is defined as any DRG in MDC 15, an admission type of newborn (ATYPE=4), or an ICD-9-CM diagnosis code for an in-hospital live-born birth.)

PSI.18 - Obstetric trauma - vaginal delivery with instrument

- This indicator flags potentially preventable trauma cases during instrument-assisted vaginal delivery. The rate is defined as the number of obstetric trauma cases (3rd or 4th degree lacerations, other obstetric lacerations) per 1,000 instrument-assisted vaginal deliveries. No risk-adjustment is made to this indicator and no significance tests performed as a result.
- The numerator refers to the number of cases of obstetric trauma on births with instrument-assisted vaginal deliveries.
- The denominator includes all vaginal delivery discharges with any listed ICD-9-CM procedure code for instrument-assisted delivery.
- Table 10 shows the number of obstetric trauma cases (obstetric trauma – vaginal delivery with instrument) among instrument-assisted vaginal delivery discharges and their corresponding observed rates, by hospital. Statewide, there were 473 cases of obstetric trauma among instrument-assisted vaginal deliveries reported in 2012 yielding a rate of 126.0 per 1,000 instrument-assisted vaginal delivery discharges.

Table 10. Obstetric trauma - vaginal delivery with instrument (per 1,000 instrument-assisted vaginal deliveries)

Hospital	# of cases	# of discharges	Obs. rate
Statewide	473	3,754	126.0
Atlanticare Regional Medical Center-Mainland	6	76	78.9
Cape Regional Medical Center	2	16	125.0 ^
Capital Health Medical Center - Hopewell	20	90	222.2
Capital Health Regional Medical Center	1	5	200.0 ^
CentraState Medical Center	5	21	238.1 ^
Chilton Memorial Hospital	21	77	272.7
Christ Hospital	1	13	76.9 ^
Clara Maass Medical Center	4	34	117.6
Community Medical Center	1	62	16.1
Cooper Hospital/University Medical Center	15	106	141.5
Englewood Hospital and Medical Center	16	103	155.3
Hackensack University Medical Center	22	344	64.0
Hackettstown Regional Medical Center	5	30	166.7
Hoboken University Medical Center	0	19	0.0 ^
Holy Name Medical Center	7	50	140.0
Hunterdon Medical Center	10	34	294.1
Jersey City Medical Center	2	104	19.2
Jersey Shore University Medical Center	11	116	94.8
JFK Medical Center/Anthony M. Yelencics	8	77	103.9
Kennedy University Hospital - Wash. Twp.	8	56	142.9
Kimball Medical Center	9	54	166.7
Meadowlands Hospital Medical Center	1	7	142.9 ^
Memorial Hospital of Salem County	0	5	0.0 ^
Monmouth Medical Center	26	216	120.4
Morristown Memorial Hospital	32	222	144.1
Mountainside Hospital	5	28	178.6 ^
Newark Beth Israel Medical Center	5	39	128.2
Newton Medical Center	4	19	210.5 ^
Ocean Medical Center - Bricktown	7	44	159.1
Our Lady of Lourdes Medical Center	5	34	147.1
Overlook Medical Center	16	119	134.5

Table 10. Obstetric trauma - vaginal delivery with instrument (per 1,000 instrument-assisted vaginal deliveries)

Hospital	# of cases	# of discharges	Obs. rate
Statewide	473	3,754	126.0
Palisades Medical Center - NY PHS	13	43	302.3
Raritan Bay Medical Center-Perth Amboy	6	30	200.0
Riverview Medical Center	9	58	155.2
RWJ University Hospital	7	38	184.2
RWJ University Hospital at Hamilton	0	15	0.0 ^
Shore Medical Center	1	11	90.9 ^
Somerset Medical Center	5	44	113.6
South Jersey Healthcare Regional MC	6	32	187.5
South Jersey Hospital-Elmer	1	5	200.0 ^
Southern Ocean Medical Center	1	8	125.0 ^
St. Barnabas Medical Center	36	277	130.0
St. Clare's Hospital-Denville	6	25	240.0 ^
St. Joseph's Hospital and Medical Center	7	33	212.1
St. Mary's Hospital (Passaic)	2	10	200.0 ^
St. Peter's University Hospital	11	179	61.5
Trinitas Regional Medical Center	6	43	139.5
UMDNJ-University Hospital	2	56	35.7
Underwood-Memorial Hospital	0	16	0.0 ^
University Medical Center at Princeton	16	85	188.2
Valley Hospital	16	149	107.4
Virtua-Memorial Hospital Burlington Cty.	12	95	126.3
Virtua-West Jersey Hospital Voorhees	35	282	124.1

Source: New Jersey 2011 UB Data.

^: Rate is based on a denominator less than 30, and should be taken with caution.

Note that this indicator is measured using observed rate only. Risk-adjustment is no more made on this indicator.

PSI.19 - Obstetric trauma - vaginal delivery without instrument

- This indicator flags cases of potentially preventable obstetric trauma during a vaginal delivery without assistance of medical instrument. The rate is defined as the number of obstetric trauma cases (3rd and 4th degree lacerations) per 1,000 vaginal deliveries that occurred without assistance of medical instrument. No risk-adjustment is made to this indicator and no significance tests performed as a result.
- The numerator includes all discharges with ICD-9-CM code for obstetric trauma in any diagnosis or procedure field (excluding instrument-assisted delivery).
- The denominator includes all vaginal delivery discharges identified by DRG or MS-DRG codes excluding cases with any listed ICD-9-CM procedure codes for instrument-assisted delivery.
- Table 11 shows the number of cases of obstetric trauma - vaginal delivery without instrument among all vaginal deliveries by hospital. Statewide, there were 1,091 cases reported for obstetric trauma - vaginal delivery without instrument in 2012. The statewide rate for this indicator is 19.1 per 1,000 vaginal deliveries.

Table 11. Obstetric trauma - vaginal deliveries without instrument (per 1,000 vaginal delivery discharges)

Hospital	# of cases	# of discharges	Observed rate
Statewide	1,091	57,223	19.1
Atlanticare Regional Medical Center-Mainland	19	1,379	13.8
Cape Regional Medical Center	7	298	23.5
Capital Health Medical Center - Hopewell	25	1,177	21.2
Capital Health Regional Medical Center	3	261	11.5
CentraState Medical Center	24	768	31.3
Chilton Memorial Hospital	18	513	35.1
Christ Hospital	3	697	4.3
Clara Maass Medical Center	12	946	12.7
Community Medical Center	14	846	16.5
Cooper Hospital/University Medical Center	33	1,491	22.1
Englewood Hospital and Medical Center	23	1,160	19.8
Hackensack University Medical Center	57	2,775	20.5
Hackettstown Regional Medical Center	5	274	18.2
Hoboken University Medical Center	17	561	30.3
Holy Name Medical Center	13	848	15.3
Hunterdon Medical Center	13	567	22.9
Jersey City Medical Center	1	776	1.3
Jersey Shore University Medical Center	16	1,127	14.2
JFK Medical Center/Anthony M. Yelensics	15	1,508	9.9
Kennedy University Hospital - Wash. Twp.	12	543	22.1
Kimball Medical Center	8	621	12.9
Meadowlands Hospital Medical Center	6	389	15.4
Memorial Hospital of Salem County	0	112	0.0
Monmouth Medical Center	44	3,244	13.6
Morristown Memmorial Hospital	48	2,241	21.4
Mountainside Hospital	17	726	23.4
Newark Beth Israel Medical Center	34	2,080	16.3
Newton Medical Center	6	319	18.8
Ocean Medical Center - Bricktown	13	615	21.1
Our Lady of Lourdes Medical Center	8	609	13.1

Table 11. Obstetric trauma - vaginal deliveries without instrument (per 1,000 vaginal delivery discharges)

Hospital	# of cases	# of discharges	Observed rate
Statewide	1,091	57,223	19.1
Overlook Medical Center	23	1,344	17.1
Palisades Medical Center - NY PHS	33	960	34.4
Raritan Bay Medical Center-Perth Amboy	17	797	21.3
Riverview Medical Center	10	686	14.6
RWJ University Hospital	45	1,367	32.9
RWJ University Hospital at Hamilton	18	713	25.2
Shore Medical Center	13	576	22.6
Somerset Medical Center	18	541	33.3
South Jersey Healthcare Regional MC	18	1,288	14.0
South Jersey Hospital-Elmer	8	239	33.5
Southern Ocean Medical Center	6	172	34.9
St. Barnabas Medical Center	77	2,856	27.0
St. Clare's Hospital-Denville	35	696	50.3
St. Joseph's Hospital and Medical Center	16	1,846	8.7
St. Mary's Hospital (Passaic)	2	461	4.3
St. Michael's Medical Center	0	2	.
St. Peter's University Hospital	53	3,317	16.0
Trinitas Regional Medical Center	27	1,453	18.6
UMDNJ-University Hospital	11	941	11.7
Underwood-Memorial Hospital	8	534	15.0
University Medical Center at Princeton	24	1,178	20.4
Valley Hospital	60	1,526	39.3
Virtua-Memorial Hospital Burlington Cty.	23	1,346	17.1
Virtua-West Jersey Hospital Voorhees	32	2,913	11.0

Source: New Jersey 2011 UB Data.

Missing (.) indicates that the denominator is less than 3 (rate is not calculated if denominator is less than 3).

Note that this indicator is measured using observed rate only. Risk-adjustment is no more made on this indicator.

Statewide PSI estimates compared to national estimates

Table 12 shows national and New Jersey statewide hospital-level patient safety indicator estimates for the 12 PSIs analyzed in this report. The national estimates are obtained from AHRQ's Comparative Data derived from the 2010 Nationwide Inpatient Sample (NIS) using PSI SAS Software (Version 4.5) while the New Jersey statewide estimates are derived from the New Jersey 2012 UB data using the same Software.

- Compared to the 2010 national PSI estimates, New Jersey has lower rates of adverse events for 7 of the 10 PSIs that are measured using rates.
- New Jersey rates were higher than the national rates only for 'Perioperative hemorrhage or hematoma', and 'Perioperative pulmonary embolism or deep vein thrombosis'.
- These differences may in part be due to differences in years of data used and/or differences in data reporting by states.

Table 12. Comparing New Jersey's Statewide PSI Rates with National Rates (per 1,000 medical and surgical discharges)

Patient Safety Indicators (PSIs)	National	New Jersey
Retained Surgical Item or Unretrieved Device Fragment Ω	930	38
Iatrogenic Pneumothorax	0.43	0.32
Postoperative Hip Fracture	0.03	0.03
Perioperative Hemorrhage or Hematoma	5.86	14.72
Perioperative Pulmonary Embolism or Deep Vein Thrombosis	4.51	5.51
Postoperative Sepsis	12.00	11.07
Postoperative Wound Dehiscence	1.85	1.67
Accidental Puncture or Laceration	2.45	1.22
Transfusion Reaction Ω	67	1
Birth Trauma - Injury to Neonate	2.10	1.73
Obstetric Trauma - Vaginal Delivery with Instrument	139.11	126.00
Obstetric Trauma - Vaginal Delivery without Instrument	22.46	19.07

National rates are from AHRQ's Comparative data for the PSIs based on the 2010 Nationwide Inpatient Sample, computed using Version 4.5 of the PSI Software while New Jersey's rates are derived from its 2012 UB data using Version 4.5 of the PSI SAS Software.

Ω : Indicator reported in volume instead of rate, because it is a rare event..

Summary of findings

This report presents occurrences of adverse events (patient safety indicators) during hospitalization in each of New Jersey hospitals. For 10 of the 12 PSIs, observed and risk-adjusted occurrence rates are provided along with confidence intervals (where applicable) to help make a statistical assessment of patient safety in New Jersey hospitals. Statewide and national estimates are also provided to help compare hospital performance to the state or to the national rates.

Comparison of a hospital's rate to the statewide rate (presented in the top row of a hospital-level PSI table) is one way to assess how well that hospital performed among its peers. A hospital's peers could be defined at many levels (e.g., teaching hospitals, urban hospitals, suburban hospitals, etc.). It is suggested that a hospital's performance be assessed by looking at its performance across the several PSI estimates presented in the 12 Tables.

According to the 2012 New Jersey data, there are substantial variations by hospital in rates of adverse events. Some hospitals exhibit significantly higher adverse event rates than the corresponding statewide rates while others have significantly lower rates.

The performances of hospitals suggested by the patient safety indicators in this report may reflect factors that do not relate to hospital performance, such as patient or physician preference, stage of illness, age, other accompanying illnesses or conditions, or the availability of specialized equipment or doctors. While the data analysis method tries to adjust for many of these factors, it is often not possible to account for all of them through statistical analysis.

Consumers should remember that doctors direct and oversee the medical care that is delivered in hospitals, prescribe tests, and prescribe medications and treatments. This report does not separate the effect of the doctor from the effect of the hospital. The quality of patient care provided in a hospital comes from how well its doctors, nurses, support staff and management work together as well as the technology and other resources available in the facility. This report is not designed to help consumers and their families choose treatment options but to help them discuss patient safety issues with their physicians.

References

AHRQ, Quality Indicators - Guide to Patient Safety Indicators:
www.qualityindicators.ahrq.gov/Downloads/Software/SAS/V31/psi_guide_v31.p

AHRQ, Quality Indicators - Patient Safety Indicators, SAS Software Documentation, Version 4.5, September 2010.

AHRQ, Quality Indicators - Patient Safety Indicators: Technical Specifications, March 2012, Version 4.5:
http://www.qualityindicators.ahrq.gov/Modules/PSI_TechSpec.aspx#

AHRQ Quality Indicator: Benchmark Data Tables for the PSI based on the 2010 Nationwide Inpatient Sample (NIS), May, 2013.
http://www.qualityindicators.ahrq.gov/Modules/psi_resources.aspx

Health Care Quality Assessment, *Patient Safety Indicators: Technical Reports*:
<http://www.nj.gov/health/healthcarequality/qi.shtml>.

For inquiries, contact the New Jersey Department of Health and Senior Services, Office of Health Care Quality Assessment, by calling (800) 418-1397, by emailing at Markos.Ezra@doh.state.nj.us by fax at (609) 984-7735.
--