

# Wildlife Populations: Peregrine Falcon

## Background

The peregrine falcon (*Falco peregrinus anatum*) bears the distinction of being the largest falcon in New Jersey, and the world's fastest bird. Like other falcons, peregrines can be distinguished by their long, pointed wings and exceptional flight speed. Adult peregrines have slate gray to bluish backs and light-colored breasts with a fine brown horizontal barring that becomes lighter with age. Immature birds are brown above with a light-colored breast that is streaked vertically with brown markings.

Superior wing speed makes the peregrine extremely proficient at catching avian prey in flight. In New Jersey, their diet consists primarily of pigeons, songbirds, shorebirds and ducks. Peregrines hunt by soaring high above their prey. Once their target is singled out, they fold their wings and drop headlong toward it. As the peregrine reaches its prey, its wings are extended in a braking motion while the legs are thrust forward in a pendulum motion. The prey is usually killed by the impact of this mid-air collision.

Historic records indicate that peregrines once nested on the cliffs above the Delaware Water Gap, and on the Palisades overlooking the Hudson River, however with the introduction and widespread use of DDT in the 1950s, peregrine populations plummeted. By 1964, peregrines had completely disappeared from the eastern half of the country, prompting federal and state governments to list the species as endangered.

With the ban of DDT in 1972, the Division of Fish and Wildlife began a restoration program in New Jersey. Young peregrines were bred in captivity by the Peregrine Fund out of Cornell University and released along the coast between 1975 and 1980. Although the coastal marshes were not considered typical nesting habitat, an abundant prey base and freedom from predation by great horned owls provided a unique opportunity to restore the peregrine population. In 1980 the first wild nesting occurred at Forsythe National Wildlife Refuge in Brigantine. Today, the peregrine's recovery in New Jersey continues at a slow, but steady pace. Each year, biologists from the Department's Endangered and Nongame Species Program (ENSP) band young falcons and gather information on nesting success and productivity. In addition, ENSP biologists continue to monitor the population to detect the presence of environmental contaminants. Since the majority of the adult population does not migrate, their health serves as a good indicator of the quality of our environment.



*Peregrine Falcon in Atlantic City (Photo by Kathleen Clark, NJDEP)*

## Status and Trends

In New Jersey today, peregrines nest along the Atlantic coast from Ocean to Cape May counties, on Delaware River bridges from Burlington to Cumberland Counties, and in natural cliff habitat in northeastern New Jersey.<sup>1</sup> Prior to 2003, all nests were on man-made structures, such as nesting towers, water towers, large bridges and high-rise buildings, however a milestone occurred in 2003 when a pair was confirmed nesting on the cliffs of the Palisades overlooking the Hudson River.

Since 1980, the peregrine population has been steadily increasing from 2 nesting pairs to a high of 37 nesting pairs in 2018 (See Figure 1). In 2006, the peregrine falcon population declined slightly to 18 pairs, but rebounded to 20 pairs in 2007. The population has since continued to grow to 37 nesting pairs in 2018 which produced 45 young for a productivity rate of 2.25 young per active nest. The natural cliff population was documented at six pairs during 2018; in the rest of the state, 20 pairs nested on towers and buildings and the remaining 11 pairs nested on bridges.

The most-watched nesting pair however resides atop 101 Hudson Street in Jersey City, where a webcam allows for remote monitoring of nesting.<sup>2</sup>




Figure 1. Number of Peregrine Falcon Nests from 1980 to 2018 in New Jersey

## Outlook and Implications

The restoration of the peregrine population in New Jersey marks an important conservation achievement. The tower and building nest sites are the consistent center of the population in New Jersey, without which the population would fluctuate widely year to year.<sup>1</sup> Offspring from New Jersey falcons have also provided birds for reintroductions in other states, thereby accelerating the recovery process in the east.

Nest success at cliff sites improved from a poor productivity rate of 0.33 in 2017, to a rate of 2.00 in 2018. Four nests were successful in producing 12 young. While 2018 was a very successful year for cliff-nesting falcons, the highly variable nest

success at the cliff territories continues to be a problem if the occupancy of historic habitat is considered important to a fully recovered population.<sup>2</sup>

Environmental contaminants such as pesticides, PCBs and heavy metals continue to threaten sensitive components of our ecosystem, including the peregrine falcon. A 2009 analysis of peregrine eggs from Mid-Atlantic States showed that southern New Jersey peregrines continued to have higher levels of organochlorines and mercury than other regions, even those in similar salt marsh habitats.<sup>3</sup> Monitoring these contaminants as well as additional ones (e.g., fire retardant chemicals called polybrominated diphenyl ethers) is important for monitoring the health of New Jersey's environment for wildlife and people.



Peregrine Falcon in Ocean County (Photo by Anthony Ferraina through NJDEP)

## More Information

[https://www.nj.gov/dep/fgw/ensp/raptor\\_info.htm#peregrine](https://www.nj.gov/dep/fgw/ensp/raptor_info.htm#peregrine)

<https://www.nj.gov/dep/fgw/peregrinecam/jcp-perfacts.htm>

<https://www.nj.gov/dep/fgw/ensphome.htm>

## References

Unless otherwise cited, the information in this report was provided by the DEP Division of Fish and Wildlife's "Peregrine Falcon Research and Management Program in New Jersey, 2018", which can be found at [https://www.nj.gov/dep/fgw/ensp/pdf/pefa18\\_report.pdf](https://www.nj.gov/dep/fgw/ensp/pdf/pefa18_report.pdf) and the Division of Fish and Wildlife's Peregrine Facts webpage, which can be found at <https://www.nj.gov/dep/fgw/peregrinecam/jcp-perfacts.htm>

<sup>1</sup>Division of Fish and Wildlife Endangered and Nongame Species Program, "Endangered, Threatened and Rare Wildlife Conservation Projects; Interim Report for Project Year September 1, 2013-August 31, 2014", [https://www.nj.gov/dep/fgw/ensp/pdf/swgreports/swg\\_report-interim13-14.pdf](https://www.nj.gov/dep/fgw/ensp/pdf/swgreports/swg_report-interim13-14.pdf), accessed 8/28/2019.

<sup>2</sup>Conserve Wildlife, Foundation of New Jersey website; <http://www.conservewildlifenj.org/education/falconcam/>, accessed 8/28/2019.

<sup>3</sup>Clark, K. E., Y. Zhao, and C. Kane. 2009. Organochlorine Pesticides, PCBs, Dioxins, and Metals in Post-term Peregrine Falcon (*Falco peregrinus*) Eggs from the Mid-Atlantic States, 1993–1999. Archives of Environmental Contamination and Toxicology 57:174-184.