

Summary of reproductive success of New Jersey black skimmers: 2002.

Colony Location/ Name	Total # Adults	Peak # Nests (Nesting Pairs)	# Young Fledged	Productivity	Problems
East Point	2	1	0	Failed	Flooding
<i>Region 3 subtotal:</i>	2	1	0	<i>Failed</i>	
Egg Island	12	6	0	Failed	Flooding
Marshelder Island	86	40	46	High	
Barrel Island	79	15	2	Low	Flooding
Holgate	130	60	8	Low	Gull Predation
Tow Island	60	0	0	Failed	Gull Nesting
<i>Region 4 subtotal:</i>	367	121	56	<i>Low</i>	
Ocean City - North	496	4	0	Failed	Flooding
<i>Region 5 subtotal:</i>	496	4	0	<i>Failed</i>	
Strathmere Natural Area	562	246	324	High	
Strathmere Bay Island	147	31	0	Failed	Flooding
<i>Region 6 subtotal:</i>	709	277	324	<i>High</i>	
Stone Harbor Point	397	210	339	High	Flooding
Champagne Island	204	100	129	High	Flooding
Coast Guard - EECEN	11	2	0	Failed	Flooding
<i>Region 7 subtotal:</i>	612	312	468	<i>High</i>	
Totals	2186	715	848	High	
Total Colonies	12				

Failed:	0.0
Low:	< 0.5
Moderate:	≤ 0.5 through < 1.0
High:	≥ 1.0

Data provided by New Jersey Division of Fish and Wildlife - Endangered and Nongame Species Program.