

## Pumpkinseed Sunfish

### Lepomis gibbosus



### **Description:**

- bright red/orange spot with light edge on black ear flap
- deep, laterally compressed body
- brassy yellow to olive sides with abundant gold and copper flecks
- wavy blue lines on cheek and opercle of adult
- dark, chain-like vertical bands on sides of young

#### Size:

• up to 16 inches; average around 7 inches

#### Range:

- Native to Atlantic Slope drainages from New Brunswick to Edisto River, South Carolina
- Great Lakes, Hudson Bay and upper Mississippi River basins from Quebec and New York west to southeastern Manitoba and North Dakota and south to northern Kentucky and Missouri
- introduced over much of southern Canada and U.S., including some Pacific drainages


#### **Facts**:

- prefer well vegetated lakes, ponds and quiet pools of creeks and small rivers
- spawning occurs late spring to late summer
- males build and protect the nest
- females, between 2 and 5 years, produce 4000 to 7000 eggs in one season
- males breed several times (about every 11 days) throughout the season
- ♦ average lifespan is 5 to 6 years
- consume insects, insect larvae, mollusks, snails, crustaceans, small fish and leeches
- effective tool for controlling mosquito populations



# Pumpkinseed Sunfish

Lepomis gibbosus



\*pumpkinseed sunfish were only caught during the years shown