

Update on PFAS Analytical Methods

Christopher A. Impellitteri, Marc Mills, Andy Gillespie USEPA-Office of Research and Development






Disclaimer

The information in the EPA presentation has been reviewed and approved for public dissemination in accordance with U.S. Environmental Protection Agency (EPA) policy. The views expressed in this presentation are those of the author(s) and do not necessarily represent the views or policies of the Agency. Any mention of trade names or commercial products does not constitute EPA endorsement or recommendation for use.



Per- & Polyfluoroalkyl Substances (PFAS)



- A class of man-made chemicals
 - Chains of carbon (C) atoms surrounded by fluorine (F) atoms, with different terminal ends
 - Complicated chemistry thousands of different variations exist in commerce
 - Widely used in industrial processes and in consumer products
 - Some PFAS are known to be PBT:
 - Persistent in the environment
 - Bioaccumulative in organisms
 - Toxic at relatively low (ppt) levels



Methods-Sampling

Guidance to avoid cross contamination in sampling

- No teflon
- Avoid contact with clothes, materials containing PFAS (e.g. some food wrappers)
- See:
 - Sampling guidance from MI https://www.michigan.gov/documents/pfasresponse/General PFAS Sampling Guidance 634597 7.pdf
 - Interstate Technology and Regulatory Council Fact Sheet: Site Characterization Considerations, Sampling Precautions, and Laboratory Analytical Methods for PFAS pfas-1.itrcweb.org/wp-content/uploads/2018/03/pfas fact sheet site characterization 3 15 18.pdf
- PFAS Quality Assurance Plan and Data Review issues
 epa.gov/fedfac/technical-fact-sheet-perfluorooctane-sulfonate-pfos-and-perfluorooctanoic-acid-pfoa-0



Targeted vs Non-Targeted

Targeted methods are methods which are applicable to a specific defined set of known analytes

- Analytical standards exist for quantitation
- Method only 'sees' analytes on the targeted list will not measure others
- 'One and done' once the analysis is complete, can't look for other analytes


Non-targeted methods involve the use of High Resolution Mass Spectrometry (HRMS) capable of identifying all analytes in a sample, known and unknown

- Can quantitate those for which laboratory standards exist, otherwise may semiquantitate based on known, structurally similar analytes
- Can screen for lists of known suspects, can discover new/unknown analytes
- Can store the HRMS data and go back later to look for analytes which were unidentified at the time of analysis but which later become known

Non-Targeted Analysis

Explore Unknown compounds using High resolution mass spectrometry. Identify a peak in a chromatogram and to ultimately predict the identity of this unknown

↓ Mass spectrometer assigns a high resolution mass for peaks observed in the chromatogram



- **↓** Software calculates the exact number and type of atoms needed to achieve the measured mass.
- **↓** Fragmentation experiments allow determination of most likely structure:

- Using mass, formula, and structure, identity can be assigned by searching against databases of known compounds
- **↓** Compare peak to commercial to confirm identification if possible



Types of Analytical Methods

Three broad classes of methods:

- EPA Standard Methods
 - Methods which have been through a multi-lab validation following a particular rulemaking or guidance effort and are available to support Agency regulatory or guidance activities
- Research Methods
 - Methods which have been developed by an EPA ORD laboratory for research purposes; QAed and peer reviewed via publication, but not multi-lab validated, not considered EPA Standard Methods
- Developmental Methods
 - Methods which are currently undergoing research, development and testing; might become Standard Methods or Research Methods



Types of Standard Methods

Three broad categories of EPA Standard Methods related to water:


- Safe Drinking Water Act Methods
 - https://www.epa.gov/dwanalyticalmethods
- Clean Water Act Methods
 - https://www.epa.gov/cwa-methods
- SW846 Methods
 - https://www.epa.gov/hw-sw846/guidance-methods-development-and-methods-validation-resource-conservation-and-recovery-act

These are generally targeted methods for solids and water



Drinking Water Method 537: Revision I


- Update: External lab validation for additional analytes by 537
 - Perfluoro-2-propoxypropanoic acid (GenX chemical HFPO-DA, CAS 13252-13-6)
 - Potassium 9-chlorohexadecafluoro-3-oxanone-1-sulfonate (9Cl-PF3ONS, CAS 73606-19-6)
 - Potassium 11-chloroeicosafluoro-3-oxaundecane-1-sulfonate (11Cl-PF3OUdS, CAS 83329-89-9)
 - Sodium dodecafluoro-3H-4,8-dioxanonate (ADONA, CAS 958445-44-8)
- Incorporated clarifications issued in EPA Technical Advisory epa.gov/sites/production/files/2016-09/documents/pfoatechnical-advisory.pdf
- Final published method (November, 2018) epa.gov/water-research-methods
- LC/MS/MS with internal standards. Single lab lowest concentration minimum reporting levels (LCMRLs) range from 0.53-6.3 ng/L





Drinking Water Method 537: "Modified"

- Method 537 often modified by analytical laboratories for use on non-drinking water samples.
- If modifications are made that are not explicitly listed in 537 or 537 Revision 1, the method is not considered 537 by EPA.
- The most common modification is inclusion of isotope dilution.





Drinking Water Method 533

Solid phase extraction/isotope dilution method targeting PFAS <C12

- Method 537 generally performs poorly for C4 compounds (e.g. PFBA, PFBS)
- Solid phase extraction, LC/MS/MS, Isotope dilution
- Will support the fifth Unregulated Contaminant Monitoring Rule
- Released December 2019
 - https://www.epa.gov/dwanalyticalmethods/analytical-methodsdeveloped-epa-analysis-unregulated-contaminants



Drinking Water Method 533

Method 533	Both Methods	Method 537.1
1H, 1H, 2H, 2H-perfluorodecane sulfonic acid (8:2 FTS)	11-chloroeicosafluoro-3-oxaundecane-1-sulfonic acid (11Cl-PF3OUdS)	N-ethyl perfluorooctanesulfonamidoacetic acid (NEtFOSAA)
1H, 1H, 2H, 2H- perfluorohexane sulfonic acid (4:2 FTS)	9-chlorohexadecafluoro-3-oxanone-1-sulfonic acid (9Cl-PF3ONS)	N-methyl perfluorooctanesulfonamidoacetic acid (NMeFOSAA)
1H, 1H, 2H, 2H-perfluorooctane sulfonic acid (6:2 FTS)	4,8-dioxa-3H-perfluorononanoic acid (ADONA) ³	Perfluorotetradecanoic acid (PFTA)
Nonafluoro-3,6-dioxaheptanoic acid (NFDHA)	Hexafluoropropylene oxide dimer acid (HFPO-DA)	Perfluorotridecanoic acid (PFTrDA)
Perfluoro (2-ethoxyethane) sulfonic acid (PFEESA)	Perfluorodecanoic acid (PFDA)	
Perfluoro-3-methoxypropanoic acid (PFMPA)	Perfluorododecanoic acid (PFDoA)	
Perfluoro-4-methoxybutanoic acid (PFMBA)	Perfluorohexanoic acid (PFHxA)	
Perfluorobutanoic acid (PFBA)	Perfluoroundecanoic acid (PFUnA)	
Perfluoroheptanesulfonic acid (PFHpS)	Perfluorobutanesulfonic acid (PFBS)	
Perfluoropentanesulfonic acid (PFPeS)	Perfluoroheptanoic acid (PFHpA)	
Perfluoropentanoic acid (PFPeA)	Perfluorohexanesulfonic acid (PFHxS)	
	Perfluorononanoic acid (PFNA)	
	Perfluorooctanoic acid (PFOA)	
	Perfluorooctanesulfonic acid (PFOS)	



Non-Drinking Water Sample Methods: SW-846 Method 8327—Direct Injection

Non-drinking water aqueous matrices:

- Groundwater
- Surface water
- Wastewater

Find a balance among sensitivity, ease of implementation, and monitoring requirements

- Simplicity
- Robustness
- Maximizing throughput for production lab use
- Minimizing sample transfers, extractions, filter steps, chemical additions (e.g., pH adjustments)

https://www.epa.gov/hw-sw846/validated-test-method-8327-and-polyfluoroalkyl-substances-pfas-using-external-standard









Non-Drinking Water Sample Methods: SW-846 Method 8327—Direct Injection

24 PFAS (including all target analytes in EPA Method 537)

Commercially available standards ("native" and isotopically labeled)

Direct injection-EPA Region 5/Chicago Regional Lab SOP

- Similar to draft American Society for Testing and Materials (ASTM)
 Method D7979
- Multi-laboratory validation study completed in 2018
- OLEM addressing public comments (closed August 26, 2019)
- Finalize in March (?) 2020

Target Quantitation Limits: 10 nanogram/L

Associated preparation method 3512 for aqueous matrices









Non-Drinking Water Sample Methods: CWA/SW846 Method—Isotope Dilution

More complex method relative to direct injection, however will

- likely be more robust for complex matrices (e.g., wastewater influents, biosolids). Account for matrix effects (e.g., sorption) through isotopically marked standard recoveries;
- afford options to meet DoD requirements; and
- allow users to perform a deeper dive based on screening (e.g. 8327) results.

40 PFAS analytes-includes all analytes listed in 537.1, 533, and SW846 8327

Non-drinking water samples

- Surface water, groundwater, wastewater
- Landfill leachates
- Solids (soils, sediments, biosolids, tissues)





Non-Drinking Water Sample Methods: CWA/SW846 Method—Isotope Dilution

Build in flexibility

- Columns
- Elution schemes

Single laboratory validation in progress

- Collaborative effort among DoD, EPA Office of Water, EPA Office of Land and Emergency Management, and EPA ORD
- Assuming single lab validation success, multi-laboratory validation will follow over Summer 2020
- Method being developed in accordance with CWA and SWA-846 protocols for method development

Target Quantitation Limits: 1-10 nanogram/L

Matrices include:

- Wastewater (influent Landfill leachate Biosolids
 - and effluent) Soil Fish tissue
- Groundwater
- Surface water

- Sediment





PFAS Analysis Marine Waters

No EPA Approved method for PFAS in Marine Waters – ORD has a research method...

LC-MS/MS Isotope dilution method

Method Details

- Covers 24 PFAS, Commercially available standards ("native" and isotopically labeled)
- Similar to those in EPA 537.1, 533, and SW 846 8327
- Target quantitation limits <1ng/L, with extracted samples
- SPE sample concentration matrix elimination (up to 500 mL, Weak-Anion Exchange (WAX))
- Accounts for matrix effects (e.g., high ionic strength and DOM)
- Adapted to estuarine sediments and TOP assay



PFAS Analysis in Fish Tissue

No EPA Approved method for PFAS in fish tissue – EPA uses commercial laboratories' proprietary methods

- LC-MS/MS with solid phase extraction and isotope dilution
- Similar to DW 533 and SWA 1600 but it is NOT a Modified Method...

Method Details

- Covers 13 carboxylic and sulfonic acids from C4 to C12, plus PFOSA; now 33 analytes
- Quantitation limits ranged from 0.25 to 1.25 ng/g (ppb) for the 13 (0.38-4.09 ng/g for 33)
- Spike stable isotopically labeled PFAS analogs into 1-2 g fillet tissue sample
- Sample digested with caustic (KOH or NaOH) methanol solution to release PFAS from tissue
- Solids removed by centrifugation, aqueous solution processed by SPE extraction
- Injected into LC-MS/MS for analysis



Longer chain PFAS C8+ most consistently present



Total Organofluorine Analysis using Combustion Ion Chromatography (TOF)

Adsorption of PFAS on to activated carbon / other sorbents



- No specialized or costly instrumentation required
- Applied for aqueous matrices and blood samples
- Removing the background inorganic F⁻ from the sample is important to make sure that the reported F⁻ is organic
- Can be developed on a wide commercial scale
- High priority for EPA. ORD working with OW-OST to develop a draft method in 2020



Total Organic Precursors (TOP)

- Developed by Houtz et al. No multi-laboratory validated standard methods.
- Available from some contract laboratories
- Does not identify individual precursor compounds

Heated oxidative conversion

60 mM Persulfate, 125 mM Base

Oxidizable PFAA Precursors

PFCAs

85 °C for 6 Hrs

Total Oxidizable Precursors = PFCAs after oxidation - PFCAs before oxidation

- Applicable for aqueous and solid matrices
- Conservative estimate of the total concentration of PFAA precursors
- More expensive; sample needs to be analyzed twice for PFAS



Summary: EPA PFAS Methods, January 2020

EPA has validated Standard Methods complete or in development for PFAS in water

- Final SDWA Methods 533 and 537.1 for available for drinking water (29 PFAS)
- Method SW846-8327 validated and undergoing final review for non-potable water (24 PFAS)
- Method CWA-1600 undergoing single and multi lab validation for non-potable water and solids (40 PFAS)

EPA has or is developing additional methods for partner use

- Fish Tissue Isotope dilution method for 13 PFAS has been used in national surveys
- Serum Isotope dilution method (targeted and non-targeted) used in biomonitoring
- Ambient air and emissions Sampling and analysis methods undergoing development and testing
- Total Organic Precursors (TOP) identify total PFAS load which may degrade to most persistent PFAS
- Total Organic Fluorine (TOF) potential rapid screening tool to identify total PFAS presence/absence
- **High resolution mass spectrometry** Continued development and application of HRMS methods for discovery of novel PFAS, suspect screening, and non-targeted analysis



For More Information

Chris Impellitteri, Ph. D.

Lead, ORD PFAS Method Development US EPA Office of Research and Development

Impellitteri.christopher@epa.gov (513) 487-2872

The views expressed in this presentation are those of the individual author and do not necessarily reflect the views and policies of the US EPA

