

When Can I Get My COVID-19 BOOSTER DOSE?*

If You Completed a COVID-19 Primary Vaccine Series**

- Get boosted 5 months after completing Pfizer or Moderna series
- Get boosted 2 months after receiving Johnson & Johnson/Janssen




If You Receive or Want to Receive Additional Vaccines

- You can get a COVID-19 vaccine/booster **before**, at the **same time**, or **after** getting other vaccines



If You Tested Positive or were Exposed to Someone with COVID-19

- Get vaccinated/boosted after isolation or quarantine period ends and symptoms have improved



Find a COVID-19 vaccine or booster:
covid19.nj.gov/finder

Questions? Call 855-568-0545



*Tell your doctor if you were diagnosed with [multisystem inflammatory syndrome](#) in children (MIS-C) or adults (MIS-A), since this can also affect the timing of your booster.

**[People with weakened immune systems](#) are recommended to receive an additional dose of an mRNA (Pfizer-BioNTech or Moderna) vaccine plus a booster dose. The timing between the additional dose and the booster depends on the vaccine you received for your primary series (starting dose(s):

If you received Pfizer-BioNTech or Moderna two-dose primary series: Booster dose is recommended at least 3 months after the additional dose for a total of four doses.

If you received a Johnson & Johnson (J&J)/Janssen one-dose primary series: Booster dose is recommended at least 2 months after the additional dose for at total of three doses.