




# **Behavioral and Clinical Characteristics of Adults with Diagnosed HIV Infection**

**Medical Monitoring Project  
New Jersey  
2015-2019 Cycle**



# Adults with Diagnosed HIV by Gender, Medical Monitoring Project, New Jersey, 2015-2019



■ *Male*


■ *Female*





■ *Transgender*

Note: Transgender persons defined as those who self-identified as transgender or who reported a gender identity different from sex assigned at birth.




# Adults with Diagnosed HIV by Sexual Orientation, Medical Monitoring Project, New Jersey, 2015-2019



 *Heterosexual or straight*       *Bisexual*  
 *Lesbian or gay*       *Other*



# Health Insurance or Coverage for Care or Antiretroviral Medications in the Past 12 Months among Adults with Diagnosed HIV, Medical Monitoring Project, New Jersey, 2015-2019




Healthcare insurance/coverage types not mutually exclusive; people could report >1 type of coverage.

\*RWHAP, Ryan White HIV/AIDS Program




# Receipt of Care and Prescription of ART Among Adults with Diagnosed HIV, Medical Monitoring Project, New Jersey, 2015-2019




\*Outpatient HIV care was defined as any documentation of the following: encounter with an HIV care provider, viral load test result, CD4 test result, HIV resistance test or tropism assay, ART prescription, PCP prophylaxis, or MAC prophylaxis

†Two elements of outpatient HIV care at least 90 days apart in each 12-month period

# Emergency Department and Hospital Admissions in the Past 12 Months Among Adults with Diagnosed HIV, Medical Monitoring Project, New Jersey, 2015-2019




# Symptoms of Depression\* in the Past 2 Weeks Among Adults With Diagnosed HIV, Medical Monitoring Project, New Jersey, 2015-2019




\*Responses to the items on the PHQ-8 were used to define "major depression" and "other depression" according to criteria from the DSM-IV. "Major depression" was defined as having at least 5 symptoms of depression; "other depression" was defined as having 2–4 symptoms of depression. The PHQ-8 classification "other depression" comprises the DSM-IV categories of dysthymia and depressive disorder, not otherwise specified, which includes minor or subthreshold depression.

# Cigarette Smoking in the Past 12 Months Among Adults with Diagnosed HIV, Medical Monitoring Project, New Jersey, 2015-2019






# Measures of Gynecologic and Reproductive Health Services in the Past 3 Years Among Women with Diagnosed HIV, Medical Monitoring Project, New Jersey, 2015-2019




# Prevention Strategies Used Among Sexually Active Adults, Medical Monitoring Project, New Jersey, 2015-2019




Note: MSM, men who have sex with men; MSW, men who have sex only with women; WSM, women who have sex with men

# Met and Unmet Needs for Non-HIV Medical Care Services in the Past 12 Months Among Adults with Diagnosed HIV, Medical Monitoring Project, New Jersey, 2015-2019



# Physical and Sexual Violence Among Adults with Diagnosed HIV, Medical Monitoring Project, New Jersey, 2015-2019



## Acknowledgements

This report is based, in part, on contributions by Medical Monitoring Project (MMP) participants, community and provider advisory boards, project area staff, interviewers, and abstractors; the Data Coordinating Center for HIV Supplemental Surveillance at ICF International; and members of the Clinical Outcomes Team, Division of HIV/AIDS Prevention, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention, CDC, Atlanta, Georgia.

