



NEW JERSEY DEPARTMENT OF HEALTH

GUIDELINES FOR THE CONTROL OF RESPIRATORY VIRUS OUTBREAKS IN LONG-TERM CARE AND OTHER INSTITUTIONAL SETTINGS

Contents

Introduction.....	2
Respiratory Outbreaks.....	2
Reporting.....	2
Case Investigation & Outbreak Investigation Steps	3
1. Confirm that an outbreak exists.....	4
2. Verify the diagnosis.....	5
3. Develop an outbreak case definition	6
4. Perform active surveillance	6
5. Document & count cases	6
6. Identify & eliminate possible transmission sources.....	7
7. Institute control measures	7
A. Maintain Standard Precautions.....	7
B. Institute Additional Transmission-Based Precautions.....	8
C. Placement & Cohorting of Patients, Staff, & Equipment	11
D. HCP Return-to-Work	11
E. Vaccinations	13
F. Antiviral Treatment & Chemoprophylaxis.....	13
G. Provide In-Service Education to All Staff on All Shifts.....	14
H. Visits from Family, Friends, & Volunteers	14
I. Environmental Measures	14
8. Evaluate the effectiveness of control measures & modify as needed	15
9. Summarize the investigation in a written report.....	15
Resources	16
References	17

Introduction

The following guidelines have been established to facilitate the investigation of viral respiratory disease outbreaks and the implementation of control measures in long-term care facilities (LTCFs) and other congregate institutional settings where healthcare is routinely delivered. For the purposes of this guidance, LTCF settings include nursing homes, skilled nursing facilities, and assisted living facilities. Throughout this guidance, the term “patient” is used and applies to individuals residing and/or being cared for at the aforementioned facilities.

New Jersey Administrative Code (N.J.A.C.), Title 8, Chapter 57 mandates that long-term care and other institutional facilities immediately report any known or suspect communicable disease outbreak by telephone to the [local health department \(LHD\)](#) with jurisdiction over the facility.¹ If the New Jersey Department of Health (NJDOH) Communicable Disease Service (CDS) receives a report of an outbreak in lieu of the LHD(s) where the outbreak is occurring, the presiding LHD will be notified by NJDOH.

Respiratory Outbreaks

Each year outbreaks of respiratory illness occur in institutional settings such as nursing homes and other LTCFs. Because of their underlying health status, patients in LTCFs are at high risk for severe illness and/or death when they become acutely ill. Historically, specific emphasis has been placed on influenza, but other respiratory viruses can also be problematic in this setting; some of these include severe acute respiratory syndrome coronavirus (SARS-CoV-2 or COVID-19), adenovirus, respiratory syncytial virus (RSV), human metapneumovirus (HMPV), rhinovirus, and human parainfluenza (HPIV).

Monitoring respiratory virus activity levels, ensuring patients and staff, including healthcare personnel (HCP), are up to date with immunizations, practicing respiratory hygiene, optimizing use of engineering controls, improving indoor air quality, and having plans for rapid testing, treatment, and/or prophylaxis are crucial in preventing respiratory outbreaks. As soon as a respiratory outbreak is suspected, the LTCF response should include laboratory testing (i.e., rapid antigen testing, PCR, and/or viral isolation) to evaluate patients and staff and determine the etiology of the outbreak.

These guidelines emphasize the following priorities regarding respiratory outbreak control:

- Early detection of an outbreak
- Stopping transmission through control measures
- Measuring morbidity and mortality
- Identifying the agent responsible for the outbreak
- Using antiviral agents to help control respiratory outbreaks

For further information on controlling viral respiratory pathogens in LTCFs, please see the [Centers for Disease Control and Prevention \(CDC\) Viral Respiratory Pathogens Toolkit for Nursing Homes](#).

Reporting

While reporting communicable disease outbreaks in healthcare institutions serves many purposes, the immediate goal is to control further spread of disease. In a residential setting, it may be difficult to determine whether an outbreak exists. An outbreak should be reported when it is suspected, and facilities should not wait for an outbreak to be confirmed before reporting.

The following are examples of confirmed or suspected outbreaks that should be reported by the facility to its LHD. This is not a comprehensive list. If you think an outbreak might be occurring (i.e., suspected), you must notify your LHD, which will collect information and provide guidance.

An outbreak may be occurring if:

1. Several patients who exhibit similar respiratory symptoms are in the same room, on the same wing of a facility, or attended a common activity; or
2. Two or more patients develop respiratory illness within 72 hours of each other; or
3. There is an increase in employee absences with staff reporting similar respiratory symptoms.

Reporting refers not only to the initial outbreak notification, but also to the provision of routine updates on the status of the outbreak. The facility and the LHD shall be in frequent contact regarding case numbers, control measures implemented, outcomes (hospitalization and/or death), and other pertinent information.

Upon receiving the initial report, the LHD shall immediately inform the NJDOH CDS of the situation.

The facility shall:

- Immediately contact the [LHD](#) by phone to report every suspected or confirmed outbreak.
- Pursuant to N.J.A.C. 8:57, a health care facility shall report incidents of infectious and communicable diseases to public health authorities. When LHD staff cannot be reached, the facility shall make the report by phone directly to NJDOH, which will then contact the LHD. Call numbers are 609-826-5964 during business hours or 609-392-2020 on nights/weekends and holidays.
- The LHD shall immediately notify NJDOH CDS of the outbreak at 609-826-5964 during business hours or 609-392-2020 after hours.

Case Investigation & Outbreak Investigation Steps

Upon notification and evaluation, NJDOH will assign an “E” number to the outbreak, which should be used for all outbreak correspondence and any laboratory sample submission to the NJ Public Health and Environmental Laboratories (PHEL).

The LHD, in consultation with the NJDOH epidemiologist, shall lead the investigation by providing the facility with guidance, support, and assistance. The LHD should consider making an on-site visit for initial evaluation and ongoing assessment.

Public health authorities and facilities will collaborate to:

1. Confirm that an outbreak exists.
2. Verify the diagnosis using clinical, epidemiological, and lab test information, considering seasonal disease occurrence.
3. Develop a case definition based on clinical and laboratory criteria.
4. Perform active surveillance.
5. Document cases in a line list.
6. Identify and eliminate transmission sources when possible.
7. Institute control measures, balancing infection control concerns with disruption of patients' quality of life routines.
8. Evaluate the effectiveness of control measures and modify as needed.
9. Summarize the investigation in a written report to communicate findings.

Note: Steps may not occur simultaneously during the course of the investigation.

1. Confirm that an outbreak exists

Gather information to confirm an outbreak is occurring within the facility; this would include initial information on the number of ill and well patients. The outbreak definitions below serve as a guide but are subject to revision based on situation-specific information. The NJDOH [COVID-19 Communicable Disease Manual](#) should be referenced for additional COVID-19 setting-specific outbreak thresholds.

Definition of a Respiratory Virus Outbreak in LTC Settings:

1. ≥ 2 facility-associated, laboratory-confirmed positive cases (e.g., influenza, RSV, adenovirus, other non-COVID-19 respiratory illnesses) identified within 72 hours of each other among patients with an epi linkage*;

OR

2. ≥ 2 facility-associated, laboratory-confirmed COVID-19 cases identified within 7 days of each other among patients with an epi linkage*;

OR

3. A sudden increase over the normal background rate of acute respiratory illness (ARI)** cases, with or without documented fever (temperature $\geq 100.4^{\circ}\text{F}$ or 2 degrees above the established baseline for that patient).

**Epi linkage is defined as having a common exposure within the facility, e.g., patients on the same unit or cared for by the same HCP. Determining epi linkages requires judgment and consulting with public health and may include weighing evidence as to whether a common exposure exists.*

***ARI is generally defined as the presence of two or more of the following symptoms: fever (subjective or objective), sore throat, cough, rhinorrhea, and nasal congestion in the absence of a known cause (e.g., seasonal allergies, COPD). Note: Elderly or medically fragile persons may manifest atypical signs of respiratory virus infection and may not present with fever.*

If an outbreak is suspected, the facility should not wait for confirmation to report to the LHD

(see “Reporting” section above).

2. Verify the diagnosis

- Determine the cause of acute respiratory illness based on the history, physical exam, and/or laboratory findings of the patient or staff member. Diagnostic testing can aid clinical judgment and guide outbreak control decisions. Be alert for noninfectious causes of symptoms such as COPD exacerbations.
- Obtain laboratory confirmation by testing specimens from several symptomatic patients and/or staff as soon as possible after illness onset. Testing exposed individuals is appropriate when clinically indicated or when testing results will inform patient/resident management or infection prevention and control decisions
 - Selection of diagnostic tests should depend on the respiratory viruses circulating in the facility and/or community. Facilities should consult with public health authorities to decide which tests should be performed.
- Lab testing in an outbreak setting may be done through the facility’s standard procedures or at PHEL with prior approval by NJDOH. The LHD or NJDOH can assist with facilitating testing at PHEL. Please refer to PHEL’s protocols on specimen collection, storage, and shipping of respiratory viruses. **All specimens sent to PHEL must be pre-approved by NJDOH and properly labeled and packaged.**
 - For detailed instructions on specimen collection, storage, packaging, labeling, and shipping of respiratory viruses, please see: [PHEL Testing, Respiratory Viruses | NJDOH](#)
- **SARS-CoV-2 Testing:**
 - **Test for SARS-CoV-2 by nucleic acid detection OR by SARS-CoV-2 antigen detection assay.** Because antigen detection assays have lower sensitivity than nucleic acid detection assays for detecting SARS-CoV-2 in upper respiratory tract specimens, a negative SARS-CoV-2 antigen detection assay result *in a symptomatic person* does not exclude SARS-CoV-2 infection and should be confirmed by either a negative result from a SARS-CoV-2 nucleic acid detection assay or a second negative antigen test result on an upper respiratory tract specimen collected 48 hours after the first negative test result. If the second antigen test is negative, per [FDA guidance](#), a third antigen test could be considered if there is a high clinical suspicion of COVID-19.
- **Influenza Testing:**
 - **The following influenza tests are recommended: molecular assays, including rapid molecular assays, other molecular tests, or reverse transcription polymerase chain reaction (RT-PCR).** If influenza molecular assays are not available, and rapid influenza diagnostic tests (RIDTs) or immunofluorescence assays are used, false negatives can occur because of lower sensitivities. If influenza is suspected and RIDTs or immunofluorescence results are negative, molecular influenza assays should be used to confirm results.
- **Other Respiratory Pathogen Testing:**

- If other respiratory pathogens other than influenza and SARS-CoV-2 are suspected, consider using multiplex RT-PCR assays targeting a panel of respiratory pathogens based on those known or suspected to be circulating, and/or based on clinical symptomatology (e.g., RSV, HPIV, and/or bacterial testing).
- After a single laboratory-confirmed case of any respiratory virus among patients has been identified, it is likely that subsequent cases of associated respiratory illness are also caused by the same organism; however, co-circulation of more than one pathogen can occur, especially when respiratory illness activity in the community is elevated. Persons developing compatible symptoms should be tested for respiratory pathogens. Ideally, at least two laboratory-confirmed cases within an incubation period are needed to confirm an outbreak's etiology. When necessary, collect additional specimens from newly ill cases. When fewer than two laboratory-confirmed cases are found, a probable infectious agent can be inferred through clinical signs and symptoms.

For additional information, please see: CDC [Viral Respiratory Pathogens Toolkit for Nursing Homes](#), CDC [Influenza virus testing in investigational outbreaks in institutional or other closed settings](#), and CDC [Overview of Testing for SARS-CoV-2](#)

3. Develop an outbreak case definition

An outbreak case definition describes the criteria that an individual must meet to be counted as an outbreak case. This includes clinical signs and symptoms, physical location, and a specific time period. Every outbreak will have a unique outbreak case definition.

The outbreak case definition will be developed by the LHD or NJDOH epidemiologist in coordination with the facility, based on the current situation. The NJDOH epidemiologist is available for consultation as needed. Outbreak cases should be counted based on this working outbreak case definition using the case line list.

4. Perform active surveillance

- Promptly identify additional cases of respiratory illness among patients and staff. Be alert for new-onset illness among exposed persons, and review patient and staff histories to identify previous cases that may not have been correctly recognized as being part of the outbreak.
- Use respiratory viral testing promptly in newly identified cases of respiratory illness so that infection control measures can be initiated to prevent further spread (e.g., antiviral prophylaxis).

5. Document & count cases

- The facility shall develop and maintain a line list. Starting and maintaining a line list helps track the progress of an outbreak. A sample facility line list for patients with acute respiratory illness may be found here: <https://www.nj.gov/health/cd/topics/outbreaks.shtml#1>.

- The facility shall share the updated line list with the LHD daily. The facility should ensure that information on hospitalizations and deaths is promptly communicated and reflected on the line list.
- The LHD shall frequently review the line list with the facility and the NJDOH epidemiologist to assess the status of the outbreak and make recommendations regarding control measures. Use the outbreak case definition when reporting and counting outbreak cases.

6. Identify & eliminate possible transmission sources

- A floor plan may be used in conjunction with the line list to document the physical locations of case-patients and ill staff to identify possible transmission routes. The facility, LHD, and NJDOH should collaborate to determine the outbreak source. Occasionally, even with thorough investigation, the source might not be identified.
- Exclude sick staff. Staff members who become sick should be sent home immediately. Please see the HCP return-to-work guidance below.
- Avoid new admissions or transfers into and out of units/wards with infected patients or facility-wide if the outbreak is widespread. Inform receiving facilities of the outbreak when transferring patients. Transfer notification applies to both ill patients and exposed well patients.
- Implement [universal masking for source control](#) for individuals residing or working in affected units or areas. Use of source control can be discontinued as a mitigation measure once the outbreak is over. During periods of higher community respiratory virus transmission, facilities should consider masking upon entry, either facility-wide or targeted to higher-risk areas or patient populations, to promote respiratory hygiene and cough etiquette.

7. Institute control measures

Evaluate the existing hierarchy of controls, including administrative (e.g., staffing, screening) and engineering controls (e.g., partitions, air quality) to minimize hazardous exposures that cannot be eliminated.

Review the following:

- OSHA [Identifying Hazard Control Options: The Hierarchy of Controls](#)
- CDC [Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings](#)

A. Maintain Standard Precautions

Standard Precautions³ are intended to be applied to the care of all patients in all healthcare settings, regardless of the suspected or confirmed presence of an infectious agent. Based on risk assessment, common-sense practices, and PPE use, Standard Precautions protect healthcare providers from injury or the spread of infection.

- Use PPE whenever possible exposure to infectious material is expected. This includes procedures and patient-care activities that are likely to generate splashes or sprays of blood, body fluids, and secretions, especially suctioning and endotracheal intubation. During aerosol-

generating procedures on patients with suspected or confirmed infections transmitted by respiratory aerosols, wear a fit-tested N95 or higher respirator in addition to gloves, gown, and face/eye protection.

- Remove gloves and gowns after each patient encounter and perform hand hygiene. Refer to the CDC Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings (2007)⁴ for examples of [Safe Donning and Removal of PPE](#).
- Perform hand hygiene immediately before touching a patient, before performing an aseptic task, before moving from work on a soiled body site to a clean body site, after touching a patient or their surroundings, after contact with blood, body fluids, or contaminated surfaces, and immediately after glove removal.
 - **Hand washing should be performed** when hands are visibly soiled, before eating, and after using the restroom.
 - **Use of an alcohol-based hand sanitizer** is the preferred method for hand hygiene in healthcare settings in most clinical situations.
 - Additional information and resources on hand hygiene can be found at:
 - [Hand Hygiene in Healthcare Settings | NJDOH](#)
 - [Guideline for Hand Hygiene in Healthcare Settings \(2002\) | CDC](#)
 - [Clinical Safety: Hand Hygiene for Healthcare Workers | CDC](#)
- Follow respiratory hygiene/cough etiquette principles. This includes providing facemasks to patients, HCP, and visitors with signs and symptoms of respiratory infection as part of Standard Precautions, which is distinct from Transmission-Based Precautions.
- Operate and maintain patient care equipment in accordance with the manufacturer's instructions for use, including cleaning and disinfection.
- Routinely clean and disinfect the environment. Handle textiles and laundry in a manner that prevents the transfer of microorganisms to others and to the environment.
- Use safe injection practices. Vaccines should be drawn up in a designated clean medication area that is not adjacent to areas where potentially contaminated items are placed. Multi-dose vials to be used for more than one patient should not be kept or accessed in the immediate patient treatment area. This is to prevent inadvertent contamination of the vial through direct or indirect contact with potentially contaminated surfaces or equipment that could then lead to infections in subsequent patients.
- Note that during times of increased respiratory illness activity, facilities may choose to implement more stringent precautions based on the facility's assessment of patient population risk, physical setup of the facility, etc.

B. Institute Additional Transmission-Based Precautions

Transmission-Based Precautions are used when the mode of transmission is not completely interrupted using Standard Precautions alone. Transmission-Based Precautions are used in addition to Standard Precautions for patients who may be infected or colonized with certain infectious agents for which

additional precautions are needed to prevent infection transmission.

Recommendations for the disease-specific type and duration of precautions are outlined in CDC Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings (2007), [Appendix A: Type and Duration of Precautions Recommended for Selected Infections and Conditions | CDC](#); Table 2. provides a list of [Clinical Syndromes or Conditions Warranting Empiric Transmission-Based Precautions in Addition to Standard Precautions](#), pending confirmatory diagnostics. Transmission-Based Precautions may be extended for immunosuppressed patients due to prolonged shedding of viral particles that may be transmitted to others.

For additional information, see [CDC Infection Control: Transmission-Based Precautions](#).

Actions Common to All Transmission-Based Precautions

In addition to Standard Precautions, some interventions/actions are to be taken in all Transmission-Based Precaution categories. These include:

- Single-patient rooms are preferred when available (and necessary for Airborne Precautions). Decisions regarding patient placement should be made on a case-by-case basis after considering infection risks to other patients in the room and available alternatives. Patients with symptoms of respiratory infection should be discouraged from using common areas when feasible.
- Limit transport and movement of patients outside of their room to medically necessary purposes and cover or contain the infected or colonized areas of the patient's body. This includes masking the patient for source control, if tolerated, for Droplet and Airborne Transmission-Based Precautions, and for respiratory pathogens as part of respiratory hygiene.
- Ensure rooms of patients on Transmission-Based Precautions are cleaned and disinfected frequently (e.g., at least daily), focusing on frequently touched surfaces and equipment in the immediate vicinity of the patient, going from the clean areas to the dirty areas.

Contact Precautions

Use Contact Precautions for patients with known or suspected infections that represent an increased risk for contact transmission. In addition to the above actions common to all Transmission-Based Precautions, for Contact Precautions, also implement the following:

- HCP should wear a gown and gloves for all interactions that may involve contact with the patient or the patient's environment. Put on a gown and gloves before room entry, and discard PPE before room exit. Do not wear the same PPE for the care of more than one patient.
- Use disposable or dedicated patient-care equipment (e.g., blood pressure cuffs, stethoscopes). If the common use of equipment for multiple patients is unavoidable, clean and disinfect such equipment before use on another patient with a U.S. Environmental Protection Agency (EPA)-registered disinfectant effective against the pathogen of concern and those commonly seen in the facility.

For an infographic (available in English and Spanish) on Contact Precautions, see: <https://www.cdc.gov/infection-control/media/pdfs/contact-precautions-sign-P.pdf>.

Droplet Precautions

Droplet Precautions are intended to prevent transmission of pathogens spread through respiratory droplets generated by coughing, sneezing, or talking. In addition to the above actions common to all Transmission-Based Precautions, for Droplet Precautions, also implement the following:

- HCP should wear a mask before entering the patient's room.
- If transport and movement of the patient outside of the room is medically necessary, instruct the patient to wear a mask as tolerated.
- **Influenza:** Droplet Precautions should be implemented for patients with suspected or confirmed influenza for seven days after illness onset or until 24 hours after the resolution of fever and respiratory symptoms, whichever is longer. People treated with influenza antiviral medications continue to shed influenza virus while on treatment. Thus, continue strict adherence to Standard Precautions (e.g., hand hygiene, respiratory hygiene, and cough etiquette) while on treatment. Refer to CDC [Infection Prevention and Control Strategies for Seasonal Influenza in Healthcare Settings](https://www.cdc.gov/infection-control/media/pdfs/droplet-precautions-sign-P.pdf).

For an infographic (available in English and Spanish) on Droplet Precautions, see: <https://www.cdc.gov/infection-control/media/pdfs/droplet-precautions-sign-P.pdf>.

Airborne Precautions

Airborne Precautions are intended to prevent the transmission of pathogens via the airborne route. In addition to the above actions common to all Transmission-Based Precautions, for Airborne Precautions, also implement the following:

- Ensure appropriate patient placement in an airborne infection isolation room (AIIR). When an AIIR is unavailable, placing the patient in a private room with the door closed and masking will reduce the likelihood of airborne transmission until the patient is transferred to a facility with an AIIR.
- HCP should wear a fit-tested NIOSH-approved N95 or higher-level respirator, which should be removed after exiting the patient care environment.
- If transport or movement outside an AIIR is necessary, instruct patients to wear a mask as tolerated. HCP transporting patients who are on Airborne Precautions do not need to wear a mask or respirator during transport if the patient is wearing a mask, unless this is recommended as part of outbreak control.

For an infographic (available in English and Spanish) on Airborne Precautions, see: <https://www.cdc.gov/infection-control/media/pdfs/airborne-precautions-sign-P.pdf>.

COVID-19 Precautions

- HCP should wear a fit-tested NIOSH-approved N95 or higher-level respirator, gown, gloves, and eye protection (i.e., goggles or a face shield that covers the front and sides of the face).
- If transport and movement of the patient outside of the room is medically necessary, instruct the patient to wear a mask as tolerated.

- Recommended duration of Transmission-Based Precautions for patients with suspected or confirmed COVID-19 can range from 10 to 20 days or more, depending on the severity of illness, immunocompromise status, and risk of other patients in the facility. Refer to [Appendix A](#) in the CDC Guideline for Isolation Precautions.

C. Placement & Cohorting of Patients, Staff, & Equipment

The LHD, in consultation with NJDOH, shall provide recommendations and guidance to the facility regarding outbreak control measures. Control measures can negatively impact patients' quality of life by restricting their lifestyle for a period of time, and staffing limitations may necessitate modification of certain control measures. Nevertheless, the facility should make every effort to institute and maintain adequate control measures until the outbreak is declared over. Additional outbreak control measures are listed below:

- As noted above, single-patient rooms are preferred when there is a concern about transmission of an infectious agent; when single rooms are limited, priority should be given to patients at higher risk of transmitting infectious material to others (e.g., uncontained secretions) and those at increased risk of infection and adverse outcomes, such as those with immunosuppression, prolonged length of stay, or total dependence on HCP for activities of daily living.
- Cohorting patients involves grouping individuals infected with the same organism together, confining their care to one area, and preventing contact with other patients. Cohorts are created based on clinical diagnosis, etiologic organism confirmation when available, epidemiology, and the mode of transmission of the infectious agent.
- In general, symptomatic patients should remain in their assigned room as much as possible, including restricting them from common areas and group activities. Medically necessary services and activities should occur in their room during the infectious isolation period.
- Assigning or cohorting HCP and equipment to specific units or patients infected with a single pathogen may help limit further transmission after implementing routine infection prevention and control measures. Staff and HCP assigned to affected unit(s) should not rotate to unaffected units until the LHD and NJDOH have determined that the outbreak is under control. This restriction includes prohibiting staff and HCP from working on unaffected units after completing their usual shift on the affected unit(s), when possible. If equipment must be shared, it should be terminally cleaned and disinfected before transporting to another unit.
- Consider modifying the use of shared care areas, such as physical therapy rooms. Modifications may include limiting the number of patients using the space at one time, such as adjusting scheduling to limit crowding.
- Consider closing the facility to new admissions if the physical setup does not allow for separation and safe care delivery within distinct cohorts.

For reference, please see: [2007 Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings](#) and CDC [Standard Precautions for All Patient Care](#).

D. HCP Return-to-Work

The following Return-to-Work Guidance is intended to help healthcare facilities protect patients, staff,

and visitors by minimizing the risk of respiratory illness transmission in the workplace. Facilities should have sick leave and return-to-work policies that are non-punitive, flexible, and aligned with public health recommendations. The following outlines criteria for when HCP who are confirmed or suspected to have or have been exposed to common respiratory viruses—such as SARS-CoV-2, seasonal influenza, or respiratory syncytial virus (RSV)—may safely resume their duties. This guidance reflects current public health recommendations and emphasizes the importance of balancing workforce capacity with infection prevention and control measures.

For this HCP return-to-work guidance, **a suspected viral respiratory infection is defined as having at least two (2) of the following symptoms**, regardless of viral test result: fever*(subjective or objective), cough, runny nose or nasal congestion, or sore throat.

HCP who are confirmed or suspected of having a viral respiratory infection should be restricted from work until:

- At least 3 days have passed since symptom onset or first positive respiratory viral test (if asymptomatic). The first day of symptom onset or the day of a positive viral test is considered day 0, where the earliest return is on day 4, AND
- Fever-free for 24 hours, without the use of antipyretics, AND
- Symptoms are improving, AND
- Feel well enough to return to work

Upon return, wear source control at least through day 7 of illness onset or first positive respiratory viral test.

Asymptomatic HCP with a known or suspected exposure to a respiratory virus do not require work restrictions; however, HCP **must wear source control at least through day 5 after the last exposure**, and **monitor for signs or symptoms for 5 days following the last exposure** (HCP who become symptomatic or test positive should be restricted from work as described above).

*Public health recommendations that are based on the presence (or absence) of fever intend to identify whether a person's temperature is elevated beyond their norm or baseline. The use of either 100.0°F (37.8°C) or 100.4°F (38.0°C) is commonly used as a cut-off for fever, but this number can vary depending on factors such as the method of measurement and the age of the person.

- Work restrictions for HCP with viral respiratory infections who have been severely to critically ill or are immunocompromised typically include consultation with an expert in pathogen transmission/ immunosuppression (e.g., an infectious disease specialist) and consideration of pathogen-specific testing to inform ongoing transmission risk.
- For HCP returning to work who care for patients/residents at the highest risk of illness, including those who are moderate to severely immunocompromised, temporary reassignment, extended restrictions, or prolonged use of source control may be considered.
- Moderate to severely immunocompromised and high-risk persons are defined in the CDC [People with Certain Medical Conditions and COVID-19 Risk Factors](#) and [People at Increased Risk for Flu Complications](#). This list may not be exhaustive; facilities should consult with occupational health and infectious disease experts.

E. Vaccinations

Vaccinations for respiratory diseases such as COVID-19, influenza, and RSV, combined with basic infection prevention and control practices, can help prevent transmission among HCP and patients. Outbreaks of vaccine-preventable and other illnesses can still occur even when vaccine coverage among patients is high. The facility should:

- Educate eligible patients and staff on the availability of vaccines for the prevention of severe illness.
- Ensure patients and staff are up to date with the recommended influenza, COVID-19, and RSV vaccines. If eligible, influenza, COVID-19, and RSV vaccines can be [simultaneously administered](#).
- Provide information using posted materials or letters to families and visitors, encouraging them to be vaccinated. For helpful resources and materials to provide for families and visitors, please see the [CDC's Vaccine Information for Adults](#).

F. Antiviral Treatment & Chemoprophylaxis

The facility shall provide antiviral treatment and chemoprophylaxis as needed.

Antiviral treatments are available for [influenza](#)² and [COVID-19](#).

- [Influenza](#): The facility's medical director should administer antiviral treatment immediately to patients who have confirmed or suspected influenza. Antiviral treatments work best when started within the first two days of symptoms; however, these medications can still help when given after 48 hours to those who are very sick, such as those who are hospitalized or at a higher risk for complications.
- [COVID-19](#): COVID-19 antiviral treatments work best when started as soon as possible, and within 3-5 (for oral treatment) or 7 (for intravenous treatment) days of symptom onset.

Chemoprophylaxis^{5,6} using antiviral medications is a key component of influenza outbreak control in institutions that house patients at higher risk of influenza complications.

- Antiviral [chemoprophylaxis](#) is meant for patients who are not exhibiting influenza-like illness but who may be (or may have been) exposed to an ill person with influenza, to prevent transmission.
- As soon as an influenza outbreak is determined, the facility should promptly initiate antiviral chemoprophylaxis for all non-ill patients living on the same unit as patient(s) with laboratory-confirmed influenza (outbreak-affected units), regardless of vaccination status.
- Consideration may be given for extending antiviral chemoprophylaxis to patients on other unaffected units based upon other factors (e.g., unavoidable mixing of patients or HCP from affected units and unaffected units).
- CDC recommends antiviral chemoprophylaxis with oseltamivir for a minimum of two weeks and continuing for at least seven days after the last known laboratory-confirmed case was identified on affected units.

- [Postexposure prophylaxis](#) may be recommended for HCP at risk for severe disease or working in an area in a healthcare facility experiencing an influenza outbreak.

G. Provide In-Service Education to All Staff on All Shifts

- In addition to all direct caregivers employed by the facility, staff includes volunteers, private duty, contracted, or agency personnel who perform housekeeping, recreational, laundry, dietary, nursing, medical, social service, physical therapy, and administrative activities.
- Provide information on the infecting organism and its transmission, Standard and Transmission-Based Precautions, movement restrictions, environmental measures, and policies on specimen collection and submission.
- Ensure current competency assessment of infection prevention and control elements and implement routine auditing with feedback to staff to monitor for compliance. Refer to the [NJDOH Infection Prevention & Control: Observational Audit vs. Competency Assessment](#).
- Contact the [LHD](#) for fact sheets or other pertinent educational materials.

Refer to the CDC's [Viral Respiratory Pathogens Toolkit for Nursing Homes](#)

H. Visits from Family, Friends, & Volunteers

- Visitors with respiratory symptoms and those suspected of having a respiratory infection should be encouraged to postpone their visit until their symptoms resolve. However, a family member determined to visit may do so under any circumstance. For such visitors, provide a mask and instruct them to limit their visit only to their respective family members and to avoid common areas and group settings.
- Post signs to reinforce infection control measures, including recommendations for using appropriate PPE and performing hand hygiene before entering and leaving patient rooms. Signage should be eye-catching and posted at building entrances as well as outside patient rooms.
- Educate all visitors (e.g., family, friends, volunteers) on the importance of vaccination to prevent infection.
- Provide tissues and/or masks to patients and visitors who are coughing or sneezing so that they can cover their mouth and nose.
- Provide tissues and alcohol-based hand rubs in common areas and waiting rooms. Encourage visitors to properly discard used tissues in waste bins.

I. Environmental Measures

Increase routine and daily cleaning and disinfection frequency during a respiratory outbreak. In general, facilities should increase the frequency of routine and daily cleaning during respiratory disease season.

- Review the approved list of disinfectants to ensure products are EPA-registered disinfectants approved for use in healthcare settings. Evaluate the product claims and contact times (how

long a disinfectant must stay wet on a surface to be effective) to ensure the selection of products are effective against organisms of concern (e.g., SARS-CoV-2, influenza) and commonly seen healthcare pathogens. Ensure staff demonstrate knowledge about the contact times and manufacturers' instructions for use. Refer to the following resources:

- CDC [Guideline for Disinfection and Sterilization in Healthcare Facilities](#)
- EPA [Registered Disinfectants](#)
 - For COVID-19 (SARS-CoV-2), see EPA [List N](#).
 - For adenoviruses, refer to the list for norovirus, see EPA [List G](#).
 - For rare or novel viruses, see EPA [List Q](#).
- Special handling of soiled linens and dietary trays is not necessary.
- Maintain general healthcare environmental cleaning techniques. Refer to the CDC [Guidelines for Environmental Infection Control in Healthcare Facilities](#).

8. Evaluate the effectiveness of control measures & modify as needed

- Evaluate and enforce adherence to infection control precautions by all staff, patients, and visitors.
- If new cases are identified after control measures have been instituted, continue outbreak control measures in consultation with the LHD and NJDOH.
- Continue control measures until no new cases are identified for two incubation periods. Generally, an outbreak is considered over when two incubation periods have passed without a new case being identified.
- When no new cases are identified after two incubation periods, outbreak control measures may be discontinued, although facilities should continue active surveillance for new cases according to LHD recommendations.

9. Summarize the investigation in a written report

The LHD shall submit a final written report to NJDOH within 30 days of completion of the investigation. See the NJDOH website for the report format, available at <https://www.nj.gov/health/forms/cds-30.pdf> (form CDS-30).

Resources

[CDC Respiratory Virus Activity Levels](#)

[NJDOH Respiratory Illness Dashboard](#)

[NJDOH Respiratory Illness Activity Reports](#)

[Information on Testing, Shipping, and Storing of Respiratory Specimens](#)

[Shipping Clinical Specimens to PHEL](#)

[How to Order Respiratory Testing Online at PHEL](#)

[CDC Viral Respiratory Pathogens Toolkit for Nursing Homes](#)

[CDC Reminder: Infection Control Actions to Take During Respiratory Virus Season](#)

[CDC Prevention Strategies for Seasonal Influenza in Healthcare Settings](#)

[CDC Interim Guidance for Influenza Outbreak Management in Long-Term Care and Post-Acute Care Facilities](#)

References

1. New Jersey Administrative Code, Title 8. Department of Health, Chapter 57: Communicable Diseases. Available at <https://casetext.com/regulation/new-jersey-administrative-code/title-8-health/chapter-57-communicable-diseases/subchapter-1-reportable-communicable-diseases>. Accessed October 27, 2023.
2. Centers for Disease Control and Prevention, Seasonal Influenza: Interim Guidance for Antiviral Use. Available at <http://www.cdc.gov/flu/professionals/antivirals/index.htm>. Accessed October 27, 2023.
3. Centers for Disease Control and Prevention, Standard Precautions for All Patient Care. Available at <https://www.cdc.gov/infectioncontrol/basics/standard-precautions.html>. Accessed May 17, 2024.
4. Centers for Disease Control and Prevention, 2007 Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings. Available at <http://www.cdc.gov/hicpac/2007IP/2007isolationPrecautions.html>. Accessed October 27, 2023.
5. Centers for Disease Control and Prevention, Interim Guidance for Influenza Outbreak Management in Long-Term Care and Post-Acute Care Facilities. Available at <https://www.cdc.gov/flu/hcp/infection-control/ltc-facility-guidance.htm>. Accessed September 18, 2025.
6. Timothy M Uyeki, Henry H Bernstein, John S Bradley, Janet A Englund, Thomas M File, Alicia M Fry, Stefan Gravenstein, Frederick G Hayden, Scott A Harper, Jon Mark Hirshon, Michael G Ison, B Lynn Johnston, Shandra L Knight, Allison McGeer, Laura E Riley, Cameron R Wolfe, Paul E Alexander, Andrew T Pavia, *Clinical Practice Guidelines by the Infectious Diseases Society of America: 2018 Update on Diagnosis, Treatment, Chemoprophylaxis, and Institutional Outbreak Management of Seasonal Influenza*, *Clinical Infectious Diseases*, Volume 68, Issue 6, 15 March 2019, Pages e1–e47, <https://doi.org/10.1093/cid/ciy866>. Accessed February 6, 2024.