

Protect Your Baby

from Respiratory Syncytial Virus (RSV)

RSV is a common virus that can cause a serious lung infection in babies. Because their lungs and airways are still developing, some babies become very sick and need to go to the hospital



TWO WAYS YOU CAN PROTECT YOUR BABY

RSV Vaccine (*Abrysvo*) During Pregnancy

- Get the RSV vaccine (Abrysvo) at 32–36 weeks of pregnancy from September 1 through March 1 to pass protection to your baby before birth.
- Helps protect your baby during their first months of life.

OR

RSV Monoclonal Antibody (*Beyfortus or Enflonsia*) After Delivery

- If you do not get the RSV vaccine during pregnancy, your baby can get a monoclonal antibody (preventive medicine) after birth.
- Babies younger than 8 months get 1 dose to help protect them during their first RSV season from October 1 through March 31.
- Some children at higher risk may need a second dose before their second RSV season (Beyfortus only).

NJHealth New Jersey Department of Health

Talk to your doctor about the best way to protect your baby.



Protect Your Baby

from Respiratory Syncytial Virus (RSV)



RSV is a common virus that can cause a serious lung infection in babies. Because their lungs and airways are still developing, some babies become very sick and need to go to the hospital

TWO WAYS YOU CAN PROTECT YOUR BABY

New Jersey Department of Health Vaccines

nj.gov/health/vaccines/

American Academy of Pediatrics (AAP)

bit.ly/4ySTxuw

Links to external sites are for informational purposes only. NJDOH does not endorse or take responsibility for the content, accuracy, or views expressed on third-party sites.

NJHealth New Jersey Department of Health

Talk to your doctor about the best way to protect your baby.