


OHI#9: Hospitalizations from or with Pneumoconiosis

Annual Age-Standardized, Rate* of Inpatient Hospitalizations from or with Pneumoconiosis, New Jersey and United States, 2000-2017



*Age-standardized rate per million residents

Estimated Annual Number and Age-Standardized Rate* of Inpatient Hospitalizations from or with Pneumoconiosis, New Jersey and United States, 2000-2017

Year	Number		Rate*	
	NJ	US	NJ	US
2000	2,101	31,755	303.3	146.5
2001	1,985	25,710	286.6	116.2
2002	1,596	32,795	230.6	151.1
2003	1,628	19,667	230.6	90.4
2004	1,539	27,146	213.5	125.0
2005	1,610	26,188	232.3	108.9
2006	1,824	20,799	250.0	86.1
2007	1,893	19,037	273.0	78.5
2008	1,910	19,097	274.0	78.9
2009	1,698	16,481	242.0	67.5
2010	1,518	10,262	213.5	41.7
2011	1,360	N/A	190.4	N/A
2012	1,314	N/A	182.6	N/A
2013	1,150	N/A	159.0	N/A
2014	1,099	N/A	150.9	N/A
2015	831	N/A	113.1	N/A
2016	624	N/A	85.3	N/A
2017	835	N/A	113.1	N/A

*Age-standardized rate per million residents

N/A: not available

Note: Caution should be taken when comparing data to other states or with national estimates due to workers' compensation (WC) eligibility criteria and availability of data from WC programs varying among states. Effective October 1, 2015 healthcare organizations and providers were required to start using ICD-10-CM codes. The transition in classification from ICD-9-CM to ICD-10-CM may not be a one-to-one match, therefore caution should be taken when comparing data before and after the transition.

Data Source: NJ Department of Health inpatient hospital discharge data; US Census Bureau, American Fact Finder [https://factfinder.census.gov/faces/nav/jsf/pages/index.xhtml?_ts=440355717799]; US 2000 Standard Population [<https://www.cdc.gov/nchs/data/statnt/statnt20.pdf>]