


OHI #9: Hospitalizations from or with Pneumoconiosis

Annual Age-Standardized, Rate* of Inpatient Hospitalizations from Asbestosis, New Jersey and United States, 2000-2017



*Age-Standardized rate per million residents

Estimated Annual Number and Age-Standardized Rate* of Inpatient Hospitalizations from Asbestosis, New Jersey and United States, 2000-2017

Year	Number		Rate*	
	NJ	US	NJ	US
2000	1,919	20,223	277.0	93.3
2001	1,779	18,523	256.7	84
2002	1,406	22,441	203.0	103.2
2003	1,437	11,862	199.9	54.6
2004	1,377	21,312	191.0	98.2
2005	1,407	14,570	202.9	60.2
2006	1,643	14,416	225.4	60.9
2007	1,720	14,841	236.0	61.7
2008	1,777	16,876	255.0	70.1
2009	1,613	11,810	214.0	48.3
2010	1,442	8,123	189.4	33.2
2011	1,306	N/A	169.7	N/A
2012	1,259	N/A	159.5	N/A
2013	1,100	N/A	137.2	N/A
2014	1,047	N/A	127.5	N/A
2015	788	N/A	93.7	N/A
2016	573	N/A	68.3	N/A
2017	784	N/A	90.1	N/A

*Age-standardized rate per million residents

N/A: not available

Note: Caution should be taken when comparing data to other states or with national estimates due to workers' compensation (WC) eligibility criteria and availability of data from WC programs varying among states. Effective October 1, 2015 healthcare organizations and providers were required to start using ICD-10-CM codes. The transition in classification from ICD-9-CM to ICD-10-CM may not be a one-to-one match, therefore caution should be taken when comparing data before and after the transition.

Data Source: NJ Department of Health inpatient hospital discharge data; US Census Bureau, American Fact Finder [https://factfinder.census.gov/faces/nav/jsf/pages/index.xhtml?_ts=440355717799]; US 2000 Standard Population [<https://www.cdc.gov/nchs/data/statnt/statnt20.pdf>]