


OHI #9: Hospitalizations from or with Pneumoconiosis

Annual Age-Standardized, Rate* of Inpatient Hospitalizations from Silicosis, New Jersey and United States, 2000-2017



*Age-Standardized rate per million residents

Estimated Annual Number and Age-Standardized Rate* of Inpatient Hospitalizations from Silicosis, New Jersey and United States, 2000-2017

Year	Number		Rate*	
	NJ	US	NJ	US
2000	47	1128	6.8	5.2
2001	43	289	6.2	1.3
2002	49	1792	7.1	8.2
2003	45	892	6.3	4.1
2004	40	1132	5.5	5.2
2005	46	754	6.7	3.1
2006	40	1033	5.5	4.1
2007	48	630	6.7	2.5
2008	43	875	6.2	3.7
2009	26	2,348	3.6	9.9
2010	34	249	4.4	1.2
2011	24	N/A	3.1	N/A
2012	23	N/A	2.7	N/A
2013	23	N/A	2.7	N/A
2014	28	N/A	3.4	N/A
2015	15	N/A	1.6	N/A
2016	26	N/A	3.2	N/A
2017	28	N/A	3.2	N/A

*Age-standardized rate per million residents

N/A: not available

Note: Caution should be taken when comparing data to other states or with national estimates due to workers’ compensation (WC) eligibility criteria and availability of data from WC programs varying among states. Effective October 1, 2015 healthcare organizations and providers were required to start using ICD-10-CM codes. The transition in classification from ICD-9-CM to ICD-10-CM may not be a one-to-one match, therefore caution should be taken when comparing data before and after the transition.

Data Source: NJ Department of Health inpatient hospital discharge data; US Census Bureau, American Fact Finder [https://factfinder.census.gov/faces/nav/jsf/pages/index.xhtml?_ts=440355717799]; US 2000 Standard Population [<https://www.cdc.gov/nchs/data/statnt/statnt20.pdf>]